



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : A61K 31/27, 31/165, 31/235, 31/265, 31/335, C07C 229/00, 233/65, 271/02, C07D 307/78, 313/04</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/25774</p> <p>(43) International Publication Date: 11 May 2000 (11.05.00)</p>
<p>(21) International Application Number: PCT/US99/25042</p> <p>(22) International Filing Date: 26 October 1999 (26.10.99)</p> <p>(30) Priority Data: 60/106,422 30 October 1998 (30.10.98) US</p> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): BAKER, Robert, K. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). KAYSER, Frank [DE/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). BAO, Jianming [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). KOTLIAR, Andrew [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). PARSONS, William, H. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). RUPPRECHT, Kathleen, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). CLAIBORNE, Christopher, F. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). LIVERTON, Nigel [GB/US]; 126 East Lincoln Avenue,</p>	<p>Rahway, NJ 07065-0907 (US). CLAREMON, David, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). THOMPSON, Wayne, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).</p> <p>(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>	
<p>(54) Title: BENZAMIDE POTASSIUM CHANNEL INHIBITORS</p> <div style="text-align: center; margin: 20px 0;"> <p style="margin-left: 400px;">(I)</p> </div> <p>(57) Abstract</p> <p>This invention relates to benzamide potassium channel inhibitors of general structural formula (I). The compounds of this invention are useful in the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and related afflictions, diseases and illnesses, and cardiac arrhythmias. Also within the scope of this invention are pharmaceutical formulations comprising a compound of formula (I) and a pharmaceutical carrier, as well as pharmaceutical formulations comprising a compound of formula (I), one or more immunosuppressive compounds and a pharmaceutical carrier.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

BENZAMIDE POTASSIUM CHANNEL INHIBITORS

BACKGROUND OF THE INVENTION

5 The present invention relates to a class of compounds that are useful as potassium channel inhibitors to treat autoimmune disorders, cardiac arrhythmias, and the like.

 Immunoregulatory abnormalities have been shown to exist in a wide variety of autoimmune and chronic inflammatory diseases, including systemic lupus erythematosus, chronic rheumatoid arthritis, 10 type I and II diabetes mellitus, inflammatory bowel disease, biliary cirrhosis, uveitis, multiple sclerosis and other disorders such as Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, psoriasis, ichthyosis, Graves ophthalmopathy and asthma.

15 Although the underlying pathogenesis of each of these conditions may be quite different, they have in common the appearance of a variety of autoantibodies and self-reactive lymphocytes. Such self-reactivity may be due, in part, to a loss of the homeostatic controls under which the normal immune system operates. Similarly, following a 20 bone-marrow or an organ transplantation, the host lymphocytes recognize the foreign tissue antigens and begin to produce antibodies which lead to graft rejection.

 One end result of an autoimmune or a rejection process is tissue destruction caused by inflammatory cells and the mediators they 25 release. Anti-inflammatory agents such as NSAID's act principally by blocking the effect or secretion of these mediators but do nothing to modify the immunologic basis of the disease. On the other hand, cytotoxic agents, such as cyclophosphamide, act in such a nonspecific fashion that both the normal and autoimmune responses are shut off. 30 Indeed, patients treated with such nonspecific immunosuppressive

agents are as likely to succumb from infection as they are from their autoimmune disease.

Cyclosporin A (CsA), which was approved by the US FDA in 1983 is currently the leading drug used to prevent rejection of
5 transplanted organs. In 1993, FK-506 (Prograf) was approved by the US FDA for the prevention of rejection in liver transplantation. CsA and FK-506 act by inhibiting the body's immune system from mobilizing its vast arsenal of natural protecting agents to reject the transplant's foreign protein. In 1994, CsA was approved by the US FDA for the
10 treatment of severe psoriasis and has been approved by European regulatory agencies for the treatment of atopic dermatitis. Though they are effective in fighting transplant rejection, CsA and FK-506 are known to cause several undesirable side effects including nephrotoxicity, neurotoxicity, and gastrointestinal discomfort. Therefore, a selective
15 immunosuppressant without these side effects still remains to be developed. Potassium channel inhibitors promise to be the solution to this problem.

The importance of potassium channels was first recognized almost fifty years ago when Hodgkin and Huxley discovered that
20 potassium ions contributed to the current that excited the squid giant. Research in the area, however, was hampered by the lack of selective, high affinity ligands for potassium channels. But the advent of recombinant DNA techniques and single cell and whole cell voltage clamp techniques has changed the slow pace of the field. Potassium
25 channels have turned out to be the most diverse family of ion channels discovered to date. They modulate a number of cellular events such as muscle contraction, neuro-endocrine secretion, frequency and duration of action potentials, electrolyte homeostasis, and resting membrane potential.

30 Potassium channels have been classified according to their biophysical and pharmacological characteristics. Salient among these

are the voltage dependent potassium channels, such as K_v1 . The K_v1 class of potassium channels is further subdivided depending on the molecular sequence of the channel, for example $K_v1.1$, $K_v1.3$, $K_v1.5$.

Functional voltage-gated K^+ channels can exist as multimeric structures formed by the association of either identical or dissimilar subunits. This phenomena is thought to account for the wide diversity of K^+ channels. However, subunit compositions of native K^+ channels and the physiologic role that particular channels play are, in most cases, still unclear.

10. Membrane depolarization by $K_v1.3$ inhibition has been shown to be an effective method to prevent T-cell proliferation and therefore has applications in many autoimmune conditions. Inhibition of K^+ channels in the plasma membrane of human T-lymphocytes has been postulated to play a role in eliciting immunosuppressive responses
15 by regulating intracellular Ca^{++} homeostasis, which has been found to be important in T-cell activation.

The $K_v1.3$ voltage-gated potassium channel is found in neurons, blood cells, osteoclasts and T-lymphocytes. The Chandy and Cahalan laboratories proposed a hypothesis that blocking the $K_v1.3$
20 channel would elicit an immunosuppressant response. (Chandy *et al.*, *J. Exp. Med.* 160, 369, 1984; Decoursey *et al.*, *Nature*, 307, 465, 1984). However, the K^+ channel blockers employed in their studies were non-selective. Until research with the peptide margatoxin, a peptide found in scorpion venom, no specific inhibitor of the $K_v1.3$ channel existed to
25 test this hypothesis. Although a laboratory (Price *et al.*, *Proc. Natl. Acad. Sci. USA*, 86, 10171, 1989) showed that charybdotoxin would block $K_v1.3$ in human T cells, charybdotoxin was subsequently shown to inhibit four different K^+ channels ($K_v1.3$ and three distinct small conductance Ca^{++} activated K^+ channels) in human T-lymphocytes,

limiting the use of this toxin as a probe for the physiological role of K_v1.3 (Leonard *et al.*, *Proc. Natl. Acad. Sci. USA*, 89, 10094, 1992).

Margatoxin, on the other hand, blocks only K_v1.3 in T-cells, and has immunosuppressant activity in both *in vitro* and *in vivo* models. (Lin *et al.*, *J. Exp. Med.*, 177, 637, 1993). The therapeutic utility of this
5 compound, however, is limited by its potent toxicity. Recently, a class of compounds has been reported that may be an attractive alternative to the above-mentioned drugs, see for example U.S. Patent Nos. 5,670,504; 5,631,282; 5,696,156; 5,679,705; and 5,696,156. While addressing some of
10 the activity/toxicity problems of previous drugs, these compounds tend to be of large molecular weight and are generally produced by synthetic manipulation of a natural product, isolation of which is cumbersome and labor intensive.

The foregoing problems are solved to a large extent by the
15 practice of the invention as herein disclosed for immunosuppression.

Atrial fibrillation (AF) is the most common sustained cardiac arrhythmia in clinical practice and is likely to increase in prevalence with the aging of the population. Currently, AF affects more than 1 million Americans annually, represents over 5% of all
20 admissions for cardiovascular diseases and causes more than 80,000 strokes each year in the United States. While AF is rarely a lethal arrhythmia, it is responsible for substantial morbidity and can lead to complications such as the development of congestive heart failure or thromboembolism. Currently available Class I and Class III
25 antiarrhythmic drugs reduce the rate of recurrence of AF, but are of limited use because of a variety of potentially adverse effects including ventricular proarrhythmia. Because current therapy is inadequate and fraught with side effects, there is a clear need to develop new therapeutic approaches.

30 Antiarrhythmic agents of Class III are drugs that cause a selective prolongation of the duration of the action potential without

significant cardiac depression. Available drugs in this class are limited in number. Examples such as sotalol and amiodarone have been shown to possess interesting Class III properties (Singh B.N., Vaughan Williams E.M. "A Third Class Of Anti-Arrhythmic Action: Effects On Atrial And Ventricular Intracellular Potentials And Other Pharmacological Actions On Cardiac Muscle, of MJ 1999 and AH 3747" Br. *J. Pharmacol* 1970; 39:675-689. and Singh B.N., Vaughan Williams E. M, "The Effect Of Amiodarone, A New Anti-Anginal Drug, On Cardiac Muscle", Br *J. Pharmacol* 1970; 39:657-667.), but these are not selective Class III agents. Sotalol also possesses Class II effects which may cause cardiac depression and is contraindicated in certain susceptible patients. Amiodarone, also is not a selective Class III antiarrhythmic agent because it possesses multiple electrophysiological actions and is severely limited by side effects (Nademanee, K. "The Amiodarone Odessey". *J. Am. Coll. Cardiol.* 1992; 20:1063-1065.) Drugs of this class are expected to be effective in preventing ventricular fibrillation. Selective class III agents, by definition, are not considered to cause myocardial depression or an induction of arrhythmias due to inhibition of conduction of the action potential as seen with Class I antiarrhythmic agents.

Class III agents increase myocardial refractoriness via a prolongation of cardiac action potential duration. Theoretically, prolongation of the cardiac action potential can be achieved by enhancing inward currents (i.e. Na^+ or Ca^{2+} currents; hereinafter I_{Na} and I_{Ca} , respectively) or by reducing outward repolarizing potassium (K^+) currents. The delayed rectifier (I_{K}) K^+ current is the main outward current involved in the overall repolarization process during the action potential plateau, whereas the transient outward (I_{to}) and inward rectifier (I_{K1}) K^+ currents are responsible for the rapid initial and terminal phases of repolarization, respectively. Cellular

electrophysiologic studies have demonstrated that I_K consists of two pharmacologically and kinetically distinct K^+ current subtypes, I_{K_r} (rapidly activating and deactivating) and I_{K_s} (slowly activating and deactivating) (Sanguinetti and Jurkiewicz, Two Components Of Cardiac Delayed Rectifier K^+ Current: Differential Sensitivity To Block By Class III Antiarrhythmic Agents, *J Gen Physiol* 1990, 96:195-215). Class III antiarrhythmic agents currently in development, including d-sotalol, dofetilide (UK-68,798), almokalant (H234/09), E-4031 and methanesulfonamide-N-[1'-6-cyano-1,2,3,4-tetrahydro-2-naphthalenyl]-3,4-dihydro-4-hydroxyspiro[2H-1-benzopyran-2,4'-piperidin]-6-yl] monochloride, predominantly, if not exclusively, block I_{K_r} . Although, amiodarone is a blocker of I_{K_s} (Balser J.R. Bennett, P.B., Hondeghem, L.M. and Roden, D.M. "Suppression Of Time-Dependent Outward Current In Guinea Pig Ventricular Myocytes: Actions Of Quinidine And Amiodarone. *Circ. Res.* 1991, 69:519-529), it also blocks I_{Na} and I_{Ca} , effects thyroid function, is as a nonspecific adrenergic blocker, and acts as an inhibitor of the enzyme phospholipase (Nademanee, K. "The Amiodarone Odyssey". *J. Am. Coll. Cardiol.* 1992; 20:1063-1065). Therefore, its method of treating arrhythmia is uncertain. Most Class III agents that are known to be in development predominantly block I_{K_r} .

Reentrant excitation (reentry) has been shown to be a prominent mechanism underlying supraventricular arrhythmias in man. Reentrant excitation requires a critical balance between slow conduction velocity and sufficiently brief refractory periods to allow for the initiation and maintenance of multiple reentry circuits to coexist simultaneously and sustain AF. Increasing myocardial refractoriness by prolonging action potential duration (APD), prevents and/or terminates reentrant arrhythmias. Most selective Class III antiarrhythmic agents currently in development, such as d-sotalol and

dofetilide predominantly, if not exclusively, block I_{kr} , the rapidly activating component of I_K found both in atrium and ventricle in man.

Since these I_{kr} blockers increase APD and refractoriness both in atria and ventricle without affecting conduction per se, 5 theoretically they represent potential useful agents for the treatment of arrhythmias like AF. These agents have a liability in that they have an enhanced risk of proarrhythmia at slow heart rates. For example, torsades de points has been observed when these compounds are utilized (Roden, D.M. "Current Status of Class III Antiarrhythmic Drug 10 Therapy", *Am J. Cardiol*, 1993; 72:44B-49B). This exaggerated effect at slow heart rates has been termed "reverse frequency-dependence", and is in contrast to frequency-independent or frequency-dependent actions (Hondegheem, L.M. "Development of Class III Antiarrhythmic Agents". *J. Cardiovasc. Cardiol.* 20 (Suppl. 2):S17-S22).

15 The slowly activating component of the delayed rectifier (I_{ks}) potentially overcomes some of the limitations of I_{kr} blockers associated with ventricular arrhythmias. Because of its slow activation kinetics however, the role of I_{ks} in atrial repolarization may be limited due to the relatively short APD of the atrium. Consequently, although I_{ks} blockers 20 may provide distinct advantage in the case of ventricular arrhythmias, their ability to affect SVT is considered to be minimal.

The ultra-rapidly activating delayed rectifier K^+ current (I_{kur}) is believed to represent the native counterpart to a cloned 25 potassium channel designated Kv1.5 and, while present in human atrium, it appears to be absent in human ventricle. Furthermore, because of its rapidity of activation and limited slow inactivation, I_{kur} is believed to contribute significantly to repolarization in human atrium. Consequently, a specific blocker of I_{kur} , that is a compound which blocks Kv1.5, would overcome the short coming of other compounds by


prolonging refractoriness by retarding repolarization in the human atrium without causing the delays in ventricular repolarization that underlie arrhythmogenic afterdepolarizations and acquired long QT syndrome observed during treatment with current Class III drugs.

5 In intact human atrial myocytes an ultra-rapidly activating delayed rectifier K^+ current I_{kur} which is also known as the sustained outward current, I_{sus} or I_{so} , has been identified and this current has properties and kinetics identical to those expressed by the human K^+ channel clone (hKv1.5, HK2) when isolated from human heart and
10 stably expressed in human (HEK-293) cell lines. (Wang, Fermini and Natel, 1993, Circ Res 73:1061-1076; Fedida et al., 1993, Circ Res 73:210-216; Snyders, Tamkun and Bennet, 1993, J Gen Physiol 101:513-543) and originally cloned from rat brain (Swanson et al., 10, Neuron 4:929-939). Although various antiarrhythmic agents are now available on the
15 market, those having both satisfactory efficacy and a high margin of safety have not been obtained. For example, antiarrhythmic agents of Class I according to the classification scheme of Vaughan-Williams ("Classification Of Antiarrhythmic Drugs" In: Cardiac Arrhythmias, edited by: E. Sandoe, E. Flensted-Jensen, K. Olesen; Sweden, Astra, Sodertalje, pp449-472, 1981) which cause a selective inhibition of the
20 maximum velocity of the upstroke of the action potential (v_{max}) are inadequate for preventing ventricular fibrillation. In addition, they have problems regarding safety, namely, they cause a depression of myocardial contractility and have a tendency to induce arrhythmias due
25 to an inhibition of impulse conduction. Beta-adrenoceptor blockers and calcium antagonists which belong to Class II and IV, respectively, have a defect in that their effects are either limited to a certain type of arrhythmia or are contraindicated because of their cardiac depressant properties in certain patients with cardiovascular disease. Their safety,
30 however, is higher than that of the antiarrhythmic agents of Class I.

The method of treatment of atrial arrhythmia presented herein provides for greater safety and efficacy as well preferentially providing treatment at fast heart rates when treatment of this type is most desired.

5 SUMMARY OF THE INVENTION

This invention relates to benzamide potassium channel inhibitors of general structural Formula I.



I


10

The compounds of this invention are useful in the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and related afflictions, diseases and illnesses, and cardiac arrhythmias. Also within the scope of this invention are

15 pharmaceutical formulations comprising a compound of Formula I and a pharmaceutical carrier, as well as pharmaceutical formulations comprising a compound of Formula I, one or more immunosuppressive compounds and a pharmaceutical carrier.

DETAILED DESCRIPTION OF THE INVENTION

An embodiment of the present invention is a compound of structural Formula I:



5

I

or a pharmaceutically acceptable salt, crystal form, or hydrate, wherein:

n is: 0, 1, 2 or 3;

r is: 0 or 1;

s is: 0 or 1;

10

R¹, R², R⁵, R⁶ and R⁷ are independently:

(1) halo, wherein halo is fluoro, chloro, bromo, or iodo,

(2) hydroxy,

(3) (C₁-C₆)-alkyl,

15

(4) HO(C₁-C₆)-alkyloxy,

(5) (C₁-C₄)-perfluoroalkyl,

(6) (C₂-C₆)-alkenyl,

(7) (C₂-C₆)-alkynyl,

(8) O[(C=O)O]_r(C₁-C₆)-alkyl, wherein the alkyl may be cyclic

20

or straight-chained,

(9) (C₁-C₆)-alkyl-S(O)_n-,

(10) phenyl,

(11) phenoxy,

(12) cyano,

- (13) nitro,
(14) CO₂H,
(15) CO(C₁-C₆)-alkyl,
(16) CO₂(C₁-C₆)-alkyl,
5 (17) CONR⁸R⁹,
(18) NR⁸R⁹,
(20) (C₂-C₆)-alkenyloxy,
(21) benzyloxy,
(22) hydrogen,
10 (23) OCF₃, or
(24) R¹ and R² or R⁶ and R⁷ can an be taken together when on adjacent carbons to form a fused benzo, dihydrofuranyl, furanyl, pyrrolidyl, dihydropyrrolidyl or 1,3-dioxolan group;

R³ and R⁴ are independently:

- 15 (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(2) hydroxy,
(3) HO(C₁-C₆)-alkyloxy,
(4) (C₁-C₄)-perfluoroalkyl,
(5) O(CO)CCl₃,
20 (6) (C₁-C₆)-alkyl-S(O)_n-,
(7) phenyl-(CH₂)_r-S(O)_n-,
(8) cyano,
(9) nitro,
(10) CO₂H,
25 (11) CO(C₁-C₆)-alkyl,
(12) CO₂(C₁-C₆)-alkyl,
(13) CONR⁸R⁹,
(14) NR⁸R⁹,
(15) O(CO)NR⁸R⁹,

- (16) azido,
- (17) $\text{NR}^8(\text{CO})\text{NR}^8\text{R}^9$,
- (18) hydrogen,
- (19) $(\text{C}_1\text{-C}_{10})\text{-alkyl}$, wherein alkyl includes cyclic as well as
5 acyclic groups and is unsubstituted or substituted with one
or two of the substituents selected from the group consisting
of:
- (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (b) hydroxy,
- 10 (c) oxo,
- (d) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_s(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (e) $(\text{C}_1\text{-C}_6)\text{-alkyl-S(O)}_n\text{-}$,
- (f) aryl- $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$,
- (g) cyano,
- 15 (h) nitro,
- (i) vinyl,
- (j) NR^8R^9 ,
- (k) $\text{O}(\text{CO})\text{NR}^8\text{R}^9$,
- (l) CHO ,
- 20 (m) CO_2H ,
- (n) $\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (o) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl may be substituted
with phenyl,
- (p) CONR^8R^9 ,
- 25 (q) aryl, wherein aryl is defined as phenyl or naphthyl,
unsubstituted or substituted with one, two or three of
the substituents selected from the group consisting of:
- (a') halo, as defined above,
- (b') hydroxy,

- 5 (c') (C₁-C₆)-alkyl,
(d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
(i') phenyl,
(j') phenoxy,
10 (k') cyano,
(l') nitro,
(m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
15 (q') NR⁸R⁹, and
(r) heteroaryl, wherein heteroaryl is defined as an
unsubstituted, monosubstituted, or disubstituted five
or six membered aromatic heterocycle containing
from 1 to 3 heteroatoms selected from the group
20 consisting of O, N and S and wherein the substituents
are members selected from the group consisting of:
(a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
25 (d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,

- (i') phenyl,
(j') phenoxy,
(k') cyano,
(l') nitro,
5 (m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
(q') NR⁸R⁹, and
10 (r') fused benzo or pyridyl group,
(s) heterocyclyl, wherein heterocyclyl is defined as a 3 to
7 atom cyclic, non-aromatic substituent containing
from 1 to 3 heteroatoms selected from the group
consisting of O, N, and S, said heterocycle being
15 unsubstituted or substituted with one, two or three
substituents selected from the group consisting of:
(a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
20 (d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
25 (i') phenyl,
(j') phenoxy,
(k') cyano,
(l') nitro,
(m') CO₂H,
30 (n') CO(C₁-C₆)-alkyl,

- (o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
(q') NR⁸R⁹,
(r') NR⁸CO(C₁-C₆)-alkyl,
5 (s') oxo,
(t') fused benzo, and
(u') fused pyridyl group;
(t) benzyl-S(O)_n-,
(u) O[(C=O)O_R]_s(C₂-C₆)-alkenyl,
10 (v) O[(C=O)O_R]_saryl,
(w) O[(C=O)O_R]_sheteroaryl,
(x) O(CH₂)_nheteroaryl, or
(y) O(CH₂)_naryl;
(20) (C₂-C₁₀)-alkenyl, wherein alkenyl is unsubstituted or
15 substituted with one or two of the substituents selected from
the group consisting of:
(a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(b) hydroxy,
(c) oxo,
20 (e) (C₁-C₆)-alkyl-S(O)_n-,
(f) phenyl-(C₁-C₆)-alkyloxy,
(g) cyano,
(h) nitro,
(i) NR⁸R⁹,
25 (j) CHO,
(k) CO₂H,
(l) CO(C₁-C₆)-alkyl,
(m) CO₂(C₁-C₆)-alkyl,
(n) CONR⁸R⁹,

- 5 (o) aryl, wherein aryl is as defined above,
(p) heteroaryl, wherein heteroaryl is as defined above,
(q) heterocyclyl, wherein heterocyclyl is as defined above,
(r) O[(C=O)O_r]_s(C₁-C₆)-alkyl, alkyl as defined above,
(s) O[(C=O)O_r]_s(C₂-C₆)-alkenyl, as defined above,
(t) O[(C=O)O_r]_saryl, aryl as defined above,
(u) O[(C=O)O_r]_sheteroaryl, heteroaryl as defined above,
(v) O(CH₂)_nheteroaryl, heteroaryl as defined above, and
(w) O(CH₂)_naryl, aryl as defined above;
- 10 (21) (C₂-C₁₀)-alkynyl, wherein alkynyl is unsubstituted or substituted with one or two of the substituents selected from the group consisting of:
- 15 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(b) hydroxy,
(c) oxo,
(d) (C₁-C₆)-alkyloxy,
(e) (C₁-C₆)-S(O)_n-,
(f) phenyl-(C₁-C₆)-alkyloxy,
20 (g) cyano,
(h) nitro,
(i) vinyl,
(j) NR⁸R⁹,
(k) NR⁸CO(C₁-C₆)-alkyl,
25 (l) CHO,
(m) CO₂H,
(n) CO(C₁-C₆)-alkyl,
(o) CO₂C(C₁-C₆)-alkyl,
(p) CONR⁸R⁹,

- (q) aryl, wherein aryl is as defined above,
- (r) heteroaryl, wherein heteroaryl is as defined above,
- (s) heterocyclyl, wherein heterocyclyl is as defined above,
- (t) $O[(C=O)O_R]_s(C_1-C_6)$ -alkyl, alkyl as defined above,
- 5 (u) $O[(C=O)O_R]_s(C_2-C_6)$ -alkenyl, as defined above,
- (v) $O[(C=O)O_R]_s$ aryl, aryl as defined above,
- (w) $O[(C=O)O_R]_s$ heteroaryl, heteroaryl as defined above
- (x) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above, and
- (y) $O(CH_2)_n$ aryl, aryl as defined above,
- 10 (22) $O[(C=O)O_R]_s(C_1-C_6)$ -alkyl, alkyl as defined above,
- (23) $O[(C=O)O_R]_s(C_2-C_6)$ -alkenyl, as defined above,
- (24) $O[(C=O)O_R]_s$ aryl, aryl as defined above,
- (25) $O[(C=O)O_R]_s$ heteroaryl, heteroaryl as defined above.
- (26) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above,
- 15 (27) aryl, wherein aryl is as defined above or
- (28) $O(CH_2)_n$ aryl, aryl as defined above;

R³ can also be any of the following when R⁴ is absent:

- (29) oxo,
- 20 (30) $=CH-(C_1-C_6)$ -alkyl, wherein alkyl is as defined above,
- (31) $=CH-(C_2-C_6)$ -alkenyl, wherein alkenyl is as defined above,
- (32) $=CH$ -aryl, wherein aryl is as defined above, or
- (33) $=CH_2$;

25 R⁸ and R⁹ are independently selected from the group consisting of:

- (1) hydrogen,
- (2) $[(C=O)O_R]_s$ aryl, wherein aryl is as defined above,
- (3) $[(C=O)O_R]_s(C_2-C_6)$ -alkenyl, wherein alkenyl is as defined above,

- (4) $[(C=O)O_r]_s(C_1-C_8)$ -alkyl, wherein alkyl is as defined above,
- (5) $(C=O)_rS(O)_n(C_1-C_8)$ -alkyl, wherein alkyl is as defined above,
- (6) $(C=O)_rS(O)_n$ -aryl, wherein aryl is as defined above, and
- (7) heterocyclyl, wherein heterocyclyl is defined above;

5

R¹⁰ is:

- (1) hydrogen,
- (2) $[(C=O)O_r]_s$ aryl, wherein aryl is as defined above, or
- (3) $[(C=O)O_r]_s(C_1-C_6)$ -alkyl, wherein alkyl is as defined above.

10

Another embodiment of the invention is the compound of

Formula 1, wherein R³ and R⁴ are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (2) hydroxy,
- (3) $HO(C_1-C_6)$ -alkyloxy,
- (4) (C_1-C_4) -perfluoroalkyl,
- (5) $O(CO)CCl_3$,
- (6) (C_1-C_6) -alkyl-S(O)_n-,
- (7) phenyl-(CH₂)_r-S(O)_n-,
- (8) cyano,
- (9) nitro,
- (10) CO₂H,
- (11) $CO(C_1-C_6)$ -alkyl,
- (12) $CO_2(C_1-C_6)$ -alkyl,
- (13) CONR⁸R⁹,
- (14) NR⁸R⁹,
- (15) O(CO)NR⁸R⁹,
- (16) azido,
- (17) NR⁸(CO)NR⁸R⁹,
- (18) hydrogen,

15

20

25

- (19) (C₁-C₆)-alkyl, wherein alkyl includes cyclic as well as acyclic groups and is unsubstituted or substituted with one or two of the substituents selected from the group consisting of:
- 5 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(b) hydroxy,
(c) oxo,
(d) O[(C=O)O_R]_S(C₁-C₆)-alkyl,
(e) (C₁-C₆)-alkyl-S(O)_n-,
10 (f) aryl-(C₁-C₆)-alkyloxy,
(g) cyano,
(h) nitro,
(i) vinyl,
(j) NR⁸R⁹,
15 (k) O(CO)NR⁸R⁹,
(l) CHO,
(m) CO₂H,
(n) CO(C₁-C₆)-alkyl,
(o) CO₂(C₁-C₆)-alkyl, wherein alkyl may be substituted
20 with phenyl,
(p) CONR⁸R⁹,
(q) aryl, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three of the substituents selected from the group consisting of:
- 25 (a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
(d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,

- 5 (f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
(i') phenyl,
(j') phenoxy,
(k') cyano,
(l') nitro,
(m') CO₂H,
10 (n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
(q') NR⁸R⁹, and
15 (r) heteroaryl, wherein heteroaryl is defined as an
unsubstituted, monosubstituted, or disubstituted five
or six membered aromatic heterocycle containing
from 1 to 3 heteroatoms selected from the group
consisting of O, N and S and wherein the substituents
are members selected from the group consisting of:
20 (a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
(d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
25 (g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
(i') phenyl,
(j') phenoxy,
(k') cyano,

- (l') nitro,
(m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
5 (p') CONR⁸R⁹,
(q') NR⁸R⁹, and
(r') fused benzo or pyridyl group,
(s) heterocyclyl, wherein heterocyclyl is defined as a 3 to
10 7 atom cyclic, non-aromatic substituent containing
from 1 to 3 heteroatoms selected from the group
consisting of O, N, and S, said heterocycle being
unsubstituted or substituted with one, two or three
substituents selected from the group consisting of:
(a') halo, as defined above,
15 (b') hydroxy,
(c') (C₁-C₆)-alkyl,
(d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
20 (g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
(i') phenyl,
(j') phenoxy,
(k') cyano,
25 (l') nitro,
(m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,

- 5 (q') NR^8R^9 ,
 (r') $\text{NR}^8\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 (s') oxo,
 (t') fused benzo, and
 (u') fused pyridyl group;
 (t) benzyl- $\text{S}(\text{O})_n\text{-}$,
 (u) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s(\text{C}_2\text{-C}_6)\text{-alkenyl}$,
 (v) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s\text{aryl}$,
 (w) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s\text{heteroaryl}$,
 10 (x) $\text{O}(\text{CH}_2)_n\text{heteroaryl}$, or
 (y) $\text{O}(\text{CH}_2)_n\text{aryl}$;
 (20) $(\text{C}_2\text{-C}_6)\text{-alkenyl}$, wherein alkenyl is unsubstituted or
 substituted with one or two of the substituents selected from
 the group consisting of:
 15 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (b) hydroxy,
 (c) oxo,
 (e) $(\text{C}_1\text{-C}_6)\text{-alkyl-S}(\text{O})_n\text{-}$,
 (f) phenyl- $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$,
 20 (g) cyano,
 (h) nitro,
 (i) NR^8R^9 ,
 (j) CHO,
 (k) CO_2H ,
 25 (l) $\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 (m) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 (n) CONR^8R^9 ,
 (o) aryl, wherein aryl is as defined above,
 (p) heteroaryl, wherein heteroaryl is as defined above,

- (q) heterocyclyl, wherein heterocyclyl is as defined above,
(r) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
(s) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
(t) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
5 (u) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above,
(v) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above, and
(w) $O(CH_2)_n$ aryl, aryl as defined above;
- (21) (C₂-C₆)-alkynyl, wherein alkynyl is unsubstituted or
10 substituted with one or two of the substituents selected from
the group consisting of:
(a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(b) hydroxy,
(c) oxo,
15 (d) (C₁-C₆)-alkyloxy,
(e) (C₁-C₆)-S(O)_n-,
(f) phenyl-(C₁-C₆)-alkyloxy,
(g) cyano,
(h) nitro,
20 (i) vinyl,
(j) NR⁸R⁹,
(k) NR⁸CO(C₁-C₆)-alkyl,
(l) CHO,
(m) CO₂H,
25 (n) CO(C₁-C₆)-alkyl,
(o) CO₂C(C₁-C₆)-alkyl,
(p) CONR⁸R⁹,
(q) aryl, wherein aryl is as defined above,
(r) heteroaryl, wherein heteroaryl is as defined above,

- (s) heterocyclyl, wherein heterocyclyl is as defined above,
 (t) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
 (u) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
 (v) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
 5 (w) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above
 (x) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above, and
 (y) $O(CH_2)_n$ aryl, aryl as defined above,
 (22) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
 (23) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
 10 (24) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
 (25) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above
 (26) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above,
 (27) aryl, wherein aryl is as defined above or
 (28) $O(CH_2)_n$ aryl, aryl as defined above;
 15
- R³ can also be any of the following when R⁴ is absent:
- (29) oxo,
 (30) =CH-(C₁-C₆)-alkyl, wherein alkyl is as defined above,
 (31) =CH-(C₂-C₆)-alkenyl, wherein alkenyl is as defined above,
 20 (32) =CH-aryl, wherein aryl is as defined above, or
 (33) =CH₂.

A preferred embodiment is the compound of Formula I,
 wherein R¹, R², R⁵, R⁶ and R⁷ are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 25 (2) hydroxy,
 (3) (C₁-C₆)-alkyl,
 (4) HO(C₁-C₆)-alkyloxy,
 (5) (C₁-C₄)-perfluoroalkyl,
 (6) (C₂-C₆)-alkenyl,

- (7) $O[(C=O)O_r]_s(C_1-C_6)$ -alkyl, wherein the alkyl may be cyclic
or straight-chained,
- (8) phenyl,
- (9) $CO(C_1-C_6)$ -alkyl,
- 5 (10) $CO_2(C_1-C_6)$ -alkyl,
- (11) $CONR^8R^9$,
- (12) NR^8R^9 ,
- (13) (C_2-C_6) -alkenyloxy,
- (14) benzyloxy,
- 10 (15) hydrogen,
- (16) OCF_3 ,
- (17) R^1 and R^2 or R^6 and R^7 can be taken together when on
adjacent carbons to form a fused benzo, dihydrofuranyl,
furanyl, pyrrolidyl, dihydropyrrolidyl or 1,3-dioxolan group;
- 15 R^3 and R^4 are independently:
- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (2) hydroxy,
- (3) $HO(C_1-C_6)$ -alkyloxy,
- (4) (C_1-C_4) -perfluoroalkyl,
- 20 (5) $O(CO)CCl_3$,
- (6) (C_1-C_6) -alkyl- $S(O)_n$ -,
- (7) phenyl- $(CH_2)_r-S(O)_n$ -,
- (8) cyano,
- (9) nitro,
- 25 (10) CO_2H ,
- (11) $CO(C_1-C_6)$ -alkyl,
- (12) $CO_2(C_1-C_6)$ -alkyl,
- (13) $CONR^8R^9$,
- (14) NR^8R^9 ,

- (15) $\text{O}(\text{CO})\text{NR}^8\text{R}^9$,
(16) azido,
(17) $\text{NR}^8(\text{CO})\text{NR}^8\text{R}^9$,
(18) hydrogen,
5 (19) $(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl includes cyclic as well as
acyclic groups and is unsubstituted or substituted with one
or two of the substituents selected from the group consisting
of:
10 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(b) hydroxy,
(c) oxo,
(d) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s(\text{C}_1\text{-C}_6)\text{-alkyl}$,
(e) aryl- $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$,
(f) NR^8R^9 ,
15 (g) $\text{O}(\text{CO})\text{NR}^8\text{R}^9$,
(h) CHO ,
(i) CO_2H ,
(j) $\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
(k) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl may be substituted
20 with phenyl,
(l) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkenyl}$,
(m) CONR^8R^9 ,
(n) aryl, wherein aryl is defined as phenyl or naphthyl,
unsubstituted or substituted with a substituent
25 selected from the group consisting of:
(a') halo, as defined above,
(b') hydroxy,
(c') $(\text{C}_1\text{-C}_6)\text{-alkyl}$,
(d') $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$,

- 5 (e') (C₁-C₆)-alkyl-S(O)_n;
 (f') phenyl,
 (g') phenoxy,
 (h') cyano,
 (i') CO₂H,
 (j') CO(C₁-C₆)-alkyl,
 (k') CO₂(C₁-C₆)-alkyl,
 (l') CONR⁸R⁹,
 (m') NR⁸R⁹, and
 10 (o) benzyl-S(O)_n-,
 (p) O[(C=O)O_r]_s(C₂-C₆)-alkenyl,
 (q) O[(C=O)O_r]_saryl,
 (r) O[(C=O)O_r]_sheteroaryl,
 (s) O(CH₂)_nheteroaryl, or
 15 (t) O(CH₂)_naryl;
- (20) (C₂-C₆)-alkenyl, wherein alkenyl is unsubstituted or substituted with a substituent selected from the group consisting of:
- 20 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (b) hydroxy,
 (c) oxo,
 (d) phenyl-(C₁-C₆)-alkyloxy,
 (e) NR⁸R⁹,
 25 (f) CHO,
 (g) CO₂H,
 (h) CO(C₁-C₆)-alkyl,
 (i) CO₂(C₁-C₆)-alkyl,
 (j) CONR⁸R⁹,

- 5 (k) aryl, wherein aryl is as defined above,
 (l) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
 (m) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
 (n) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above,
 (o) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above, and
 (p) $O(CH_2)_n$ aryl, aryl as defined above;
 (21) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
 (22) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
 (23) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
 10 (24) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above
 (25) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above,
 (26) aryl, wherein aryl is as defined above or
 (27) $O(CH_2)_n$ aryl, aryl as defined above;
 15 R^3 can also be any of the following when R^4 is absent:
 (28) oxo,
 (29) $=CH-(C_1-C_6)$ -alkyl, wherein alkyl is as defined above,
 (30) $=CH-(C_2-C_6)$ -alkenyl, wherein alkenyl is as defined above,
 (31) $=CH$ -aryl, wherein aryl is as defined above, or
 20 (32) $=CH_2$;

R^8 and R^9 are independently selected from the group consisting of:

- (1) hydrogen,
 (2) $[(C=O)O_R]_S$ aryl, wherein aryl is as defined above,
 25 (3) $[(C=O)O_R]_S(C_2-C_8)$ -alkenyl, wherein alkenyl is as defined
 above, and
 (4) $[(C=O)O_R]_S(C_1-C_8)$ -alkyl, wherein alkyl is as defined above,
 and

R¹⁰ is:

- (1) hydrogen, or
- (2) [(C=O)O_r]_s(C₁-C₆)-alkyl, wherein alkyl is as defined above.

Another preferred emodiment is the compound of Formula

5 I, wherein R¹, R², R⁵, R⁶ and R⁷ are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (2) hydroxy,
- (3) (C₁-C₃)-alkyl,
- (4) (C₂-C₃)-alkenyl,
- 10 (5) O(C₁-C₄)-alkyl, wherein the alkyl may be cyclic or straight-chained,
- (6) O(CO)CH₃,
- (7) CO(C₁-C₃)-alkyl,
- (8) CO₂(C₁-C₃)-alkyl,
- 15 (9) hydrogen,
- (10) R¹ and R² or R⁶ and R⁷ can an be taken together when on adjacent carbons to form a fused benzo, dihydrofuranyl, furanyl, pyrrolidyl, dihydropyrrolidyl or 1,3-dioxolan group;

20 R³ and R⁴ are independently:

- (1) hydrogen,
 - (2) (C₁-C₆)-alkyl, wherein alkyl includes cyclic as well as acyclic groups and is unsubstituted or substituted with one or two of the substituents selected from the group consisting of:
 - 25 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 - (b) hydroxy,
 - (c) oxo,
 - (d) O[(C=O)O_r]_s(C₁-C₆)-alkyl, wherein r and s are
- 30 independently 0 or 1,

- 5
- (e) $\text{CO}_2(\text{C}_2\text{-C}_3)\text{-alkenyl}$,
 - (f) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s(\text{C}_1\text{-C}_6)\text{-alkenyl}$, wherein r and s are independently 0 or 1,
 - (g) NR^8R^9 ,
 - (h) $\text{O}(\text{CO})\text{NR}^8\text{R}^9$,
 - (i) CHO ,
 - (j) CO_2H ,
 - (k) $\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 - (l) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl may be substituted with phenyl,
 - (m) CONR^8R^9 ,
 - (n) aryl, wherein aryl is defined as phenyl, unsubstituted or substituted with a substituent selected from the group consisting of:
 - 15 (a') halo, as defined above,
 - (b') hydroxy,
 - (c') $(\text{C}_1\text{-C}_6)\text{-alkyl}$, and
 - (d') $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$, and
 - (o) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s(\text{C}_2\text{-C}_6)\text{-alkenyl}$, wherein r and s are independently 0 or 1,
- 20
- (3) $(\text{C}_2\text{-C}_6)\text{-alkenyl}$,
 - (4) aryl, wherein aryl is as defined above or
 - (5) $\text{O}(\text{CH}_2)_n\text{aryl}$, wherein aryl is as defined above;
- 25

R^3 can also be any of the following when R^4 is absent:

- (6) $=\text{CH}-(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl is as defined above,
- (7) $=\text{CH}-(\text{C}_2\text{-C}_6)\text{-alkenyl}$, wherein alkenyl is as defined above,

R⁸ and R⁹ are independently selected from the group consisting of:

- (1) hydrogen,
- (2) [(C=O)O_r]_saryl, wherein aryl is as defined above and r and s are independently 0 or 1,
- 5 (3) (C₂-C₈)-alkenyl, wherein alkenyl is as defined above, and
- (4) (C₁-C₆)-alkyl, wherein alkyl is as defined above, and

R¹⁰ is:

- (1) hydrogen, or
- 10 (2) (C=O)(C₁-C₃)-alkyl, wherein alkyl is as defined above.

A most preferred emodiment is a compound which is selected from the group consisting of: 1-(2-methoxyphenyl)-1-oxo-2-aza-(S)-4-*i*-butyl-4-phenylbutane, 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-((3-allyloxycarbonyloxy)propyl))-4-phenylbutane, 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-phenylbutane, 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(phenyl)-butane, and 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-(3-hydroxypropyl)-4-phenylbutane, or a pharmaceutically acceptable salt thereof.

20 Also encompassed by the present invention is a method of treating a condition in a mammal, the treatment of which is effected or facilitated by Kv1.3 inhibition, which comprises administering the compound of Claim 1 in an amount that is effective at inhibiting Kv1.3. Preferred conditions include: resistance by

25 transplantation of organs or tissue, graft-versus-host diseases brought about by medulla ossium transplantation, rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis, juvenile-onset or recent-onset diabetes mellitus, posterior uveitis,

30 allergic encephalomyelitis, glomerulonephritis, infectious diseases

- caused by pathogenic microorganisms, inflammatory and hyperproliferative skin diseases, psoriasis, atopic dermatitis, contact dermatitis, eczematous dermatitises, seborrhoeis dermatitis, Lichen planus, Pemphigus, bullous pemphigoid, Epidermolysis
- 5 bullosa, urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus, acne, Alopecia areata, keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus,
- 10 Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, pollen allergies, reversible obstructive airway disease, bronchial asthma, allergic asthma, intrinsic asthma, extrinsic asthma, dust asthma, chronic or inveterate asthma, late asthma and airway hyper-responsiveness,
- 15 bronchitis, gastric ulcers, vascular damage caused by ischemic diseases and thrombosis, ischemic bowel diseases, inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and leukotriene B₄-mediated diseases, Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis,
- 20 Crohn's disease, ulcerative colitis, migraine, rhinitis, eczema, interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome, diabetic nephropathy, multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis, mononeuritis, radiculopathy, hyperthyroidism, Basedow's disease,
- 25 pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia, anerythroplasia, osteoporosis, sarcoidosis, fibroid lung, idiopathic interstitial pneumonia, dermatomyositis, leukoderma vulgaris,
- 30 ichthyosis vulgaris, photoallergic sensitivity, cutaneous T cell lymphoma, arteriosclerosis, atherosclerosis, aortitis syndrome,

polyarteritis nodosa, myocardosis, scleroderma, Wegener's granuloma, Sjogren's syndrome, adiposis, eosinophilic fascitis, lesions of gingiva, periodontium, alveolar bone, substantia ossea dentis, glomerulonephritis, male pattern alopecia or alopecia senilis

5 by preventing epilation or providing hair germination and/or promoting hair generation and hair growth, muscular dystrophy; Pyoderma and Sezary's syndrome, Addison's disease, ischemia-reperfusion injury of organs which occurs upon preservation, transplantation or ischemic disease, endotoxin-shock,

10 pseudomembranous colitis, colitis caused by drug or radiation, ischemic acute renal insufficiency, chronic renal insufficiency, toxinosis caused by lung-oxygen or drugs, lung cancer, pulmonary emphysema, cataracta, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring, corneal alkali burn,

15 dermatitis erythema multiforme, linear IgA ballous dermatitis and cement dermatitis, gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution, aging, carcinogenis, metastasis of carcinoma and hypobaropathy, disease caused by histamine or leukotriene-C4 release, Behcet's disease, autoimmune

20 hepatitis, primary biliary cirrhosis sclerosing cholangitis, partial liver resection, acute liver necrosis, necrosis caused by toxin, viral hepatitis, shock, or anoxia, B-virus hepatitis, non-A/non-B hepatitis, cirrhosis, alcoholic cirrhosis, hepatic failure, fulminant hepatic failure, late-onset hepatic failure, "acute-on-chronic" liver failure,

25 augention of chemotherapeutic effect, cytomegalovirus infection, HCMV infection, AIDS, cancer, senile dementia, trauma, and chronic bacterial infection. A most preferred condition is an autoimmune disease.

A further embodiment of the invention is a method of

30 preventing or treating resistance to transplantation or transplantation rejection of organs or tissues in a patient in need

thereof, which comprises administering a therapeutically effective amount of a compound of Formula I. Also encompassed within the invention is a method of suppressing the immune system in a subject in need thereof, which comprises administering to the subject an
5 immune suppressing amount of a compound of Formula 1.

Another embodiment is a pharmaceutical formulation comprising a pharmaceutically acceptable carrier and the compound of Formula 1 or a pharmaceutically acceptable crystal form or hydrate thereof. A subembodiment of the invention is a
10 pharmaceutical formulation of the compound of Formula I, comprising, in addition, a second immunosuppressive agent. Preferred immunosuppressive agents include, but are not limited to, azathioprine, brequinar sodium, deoxyspergualin, mizaribine, mycophenolic acid morpholino ester, cyclosporin, FK-506 and
15 rapamycin.

Yet another embodiment is a method of treating a condition in a mammal, the treatment of which is effected or facilitated by $K_v1.5$ inhibition, which comprises administering a compound of Formula 1 in an amount that is effective at inhibiting $K_v1.5$. A preferred embodiment
20 is a method of preventing or treating cardiac arrhythmias in a mammal, which comprises administering a therapeutically effective amount of a compound of Formula I.

The compounds of the present invention have asymmetric centers and this invention includes all of the optical isomers and
25 mixtures thereof. Unless specifically mentioned otherwise, reference to one isomer applies to both isomers.

In addition compounds with carbon-carbon double bonds may occur in Z- and E- forms with all isomeric forms of the compounds being included in the present invention.

30 As used herein, the term "alkyl", unless otherwise indicated, includes those alkyl groups of a designated number of carbon

atoms of either a straight, branched, or cyclic configuration (carbocycles). Examples of "alkyl" include methyl, ethyl, propyl, isopropyl, butyl, sec-and tert-butyl, pentyl, hexyl, heptyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, and the like.

- 5 "Alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge, such as methoxy, ethoxy, propoxy, butoxy and pentoxy. The following illustrate the foregoing definitions: "(C₁-C₃)-alkyl" may be methyl, ethyl, propyl, isopropyl, or cyclopropyl. Similarly, "O-(C₁-C₃)-alkyl" may be methoxy, ethoxy, n-propoxy, i-
- 10 propoxy, or cyclopropoxy. In some cases, a C₀ designation is used, as in "-(C₀-C₂)-alkyl-phenyl." In such a case, the substituent is intended to be any of the following: phenyl, benzyl, 1-phenylethyl, or 2-phenylethyl. In certain definitions, the alkyl may be substituted with one or more substituents. For example a definition which reads "(C₁-C₂)-alkyl,
- 15 substituted with one or two substituents selected from oxo, hydroxy, and halo" is intended to include C(O)CH₃, CH₂BrCH₃, CO₂H, C(OH)CH₃, CH₂CH₂(OH), CH₂CO₂H, CHBrCH₂Cl, CHO, and so on.

- "Alkenyl" is intended to include hydrocarbon chains of a specified number of carbon atoms of either a straight- or branched-
- 20 configuration and at least one unsaturation, which may occur at any point along the chain, such as ethenyl, propenyl, butenyl, pentenyl, dimethyl pentenyl, and the like, and includes E and Z forms, where applicable. "Halogen" and "halo", as used herein, mean fluoro, chloro, bromo and iodo.

- 25 The term "aryl," unless specifically defined otherwise, defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three of the substituents selected from the group consisting of halo, hydroxy, alkyl, perfluoroalkyl, alkenyl, alkynyl, alkyloxy, alkyl-S(O)_n-, phenyl, phenoxy, cyano, nitro, CO₂H, CO-alkyl, CO₂-alkyl,
- 30 CONR⁸R⁹, and NR⁸R⁹.

The term "heteroaryl" as utilized herein, unless specifically defined otherwise, is intended to include the following: an unsubstituted, monosubstituted, or disubstituted five or six membered aromatic heterocycle containing from 1 to 3 heteroatoms selected from the group consisting of O, N and S and wherein the substituent is halo, hydroxy, alkyl, perfluoroalkyl, alkenyl, alkynyl, alkyloxy, -alkyl-S(O)_n-, phenyl, phenoxy, cyano, nitro, CO₂H, CO-alkyl, CO₂-alkyl, CONR⁸R⁹, NR⁸R⁹, or a fused benzo or pyridyl group. Heteroaryl groups within the scope of this definition include but are not limited to: acridinyl, carbazolyl, cinnolinyl, quinoxalinyl, pyrazolyl, indolyl, benzotriazolyl, furanyl, thienyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, and pyrrolyl which are substituted or unsubstituted as defined above.

In the compounds of Formula I, the heteroaryl or aryl groups may be optionally substituted with the substituents listed above at any available carbon atom or nitrogen atom (if present), but compounds bearing certain substituents, directly substituted to a nitrogen may be relatively unstable and are not preferred. The heteroaryl may also be fused to a second 5-, 6-, or 7-membered ring containing one or two oxygens such as: dioxolanyl, dihydrofuranyl, dihydropyranyl, and dioxanyl. Disubstituted aryl groups may be ortho, para or meta and all three are intended unless specifically defined otherwise.

"Heterocyclyl" is defined as a 3 to 7 atom cyclic, non-aromatic substituent containing from 1 to 3 heteroatoms selected from the group consisting of O, N, and S, which is unsubstituted or substituted with one, two or three substituents selected from the group consisting of halo, hydroxy, alkyl, perfluoroalkyl, alkenyl, alkynyl, alkyloxy, alkyl-S(O)_n-, phenyl, phenoxy, cyano, nitro, CO₂H, COalkyl, CO₂-alkyl, CONR⁸R⁹, NR⁸R⁹, NR⁸CO-alkyl, oxo, fused benzo, and fused pyridyl group.

Pharmaceutically acceptable salts include both the metallic (inorganic) salts and organic salts; a list of which is given in *Remington's Pharmaceutical Sciences*, 17th Edition, pg. 1418 (1985). It is well known to one skilled in the art that an appropriate salt form is

5 chosen based on physical and chemical stability, flowability, hydroscopicity and solubility. As will be understood by those skilled in the art, pharmaceutically acceptable salts include, but are not limited to salts of inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide, and nitrate or salts of an organic acid such as malate,


10 maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate or palmoate, salicylate and stearate. Similarly pharmaceutically acceptable cations include, but are not limited to sodium, potassium, calcium, aluminum, lithium and ammonium (especially ammonium salts with secondary amines).

15 Preferred salts of this invention for the reasons cited above include potassium, sodium, calcium and ammonium salts. Also included within the scope of this invention are crystal forms, hydrates and solvates of the compounds of Formula I.

Methods for preparing the compounds of this invention are

20 illustrated in the following schemes and examples. Other synthetic routes would be readily apparent to those skilled in the art.

REACTION SCHEME A




Many methods are available to prepare the amine precursors that serve as starting materials for the compounds of this invention. The substituted or unsubstituted phenylacetonitrile substrates shown in Scheme A are commercially available or can be prepared by procedures well known in the art. As shown in Scheme A, the phenylacetonitrile precursors are deprotonated with bases such as potassium t-butoxide (KOtBu) or sodium hydride (NaH) in an aprotic solvent such as THF and alkylated with alkyl halides. This leads primarily to bis-alkylation. The resulting arylacetonitriles are then reduced to amines with suitable reducing agents such as excess lithium aluminum hydride ($LiAlH_4$) in an aprotic solvent such as THF, preferably at elevated temperatures.

Amine intermediates are also prepared from substituted phenylacetic acid precursors, many which are commercially available in chiral or racemic form. The acid substrate is reduced with $LiAlH_4$ in THF to give the alcohol. The hydroxy group is derivatized as the methanesulfonate with methanesulfonyl chloride in pyridine or in THF or dichloromethane with triethylamine or pyridine. This intermediate is then reacted with sodium azide in a polar aprotic solvent such as

- dimethylformamide. Alternatively, lithium azide in THF can be used. Finally, the azide is reduced to the corresponding amino group by hydrogenation using a palladium catalyst. This and other procedures are described in March, J.; *Advanced Organic Chemistry*, 4th ed., John Wiley & Sons, New York, pp. 428 and 1219 (1992).

REACTION SCHEME B




- 10 In reaction Scheme B, R^4 groups are appended by asymmetric alkylation of phenylacetyloxazolidinone imides as reviewed by Evans, *Asymmetric Synthesis*, vol. 3, Academic Press Inc., New York, p 2-101 (1984). Thus, alkylation using (R)-(-)-4-phenyl-2-oxazolidinone provides the R^4 substituent in the (S)-configuration. The

(S)-(-)-4-phenyl-2-oxazolidinone gives R^4 in the (R)-configuration. The oxazolidinone group is removed with LiAlH_4 to give the hydroxy intermediate which is then converted as described in Reaction Scheme A.

5

REACTION SCHEME C




As shown in Scheme C, the amine intermediates are then
10 acylated with acid chlorides in aprotic solvents such as THF and CH_2Cl_2
with a base such as triethylamine to give the corresponding benzamides.
The acid chlorides, when not purchased, are prepared by stirring the
carboxylic acids in reagents such as oxalyl chloride or thionyl chloride.
Alternatively, amides can be prepared by reaction of benzoic acids with
15 the amine using the standard coupling conditions as described in
March, J.; Advanced Organic Chemistry, 4th ed., John Wiley & Sons,
New York, pp. 417-424 (1992). One such coupling reagent is dicyclohexyl
carbodiimide as reviewed by Bodanski, The Practice of Peptide Synthesis,
Springer, New York, (1984).

20

25

REACTION SCHEME D




As shown in Scheme D, in an embodiment of this invention, derivatives containing olefins are hydroborated to give primary alcohols as reviewed in March, J.; Advanced Organic Chemistry, 4th ed., John Wiley & Sons, New York, pp. 783-789 (1992). For optimal regioselectivity, a substituted borane reagent such as 9-borabicyclo(3.3.1)nonane (9-BBN) is preferred. Cleavage of the borane intermediate with hydrogen peroxide provides the alcohol.

The primary alcohol is then converted to an aldehyde by a Swern oxidation procedure with oxalyl chloride in DMSO as reviewed by Mancuso and Swern, Synthesis, 165-185 (1981).


The primary alcohol or aldehyde is converted to a carboxylic acid by a variety of oxidation procedures as referenced by Larock, Comprehensive Organic Transformation, VCH, New York, pp 93-97 (1989).

REACTION SCHEME E



In Reaction Scheme E, the olefin group can be oxidatively
 5 cleaved with ozone. The ozonide can be reduced with sodium
 borohydride to give the corresponding alcohol, or it can be decomposed to
 the corresponding aldehyde as described in March, J.; Advanced
 Organic Chemistry, 4th ed., John Wiley & Sons, New York, pp. 1177-1181
 (1992). Alternatively, the aldehyde is prepared by cleavage with catalytic
 10 osmium tetroxide and periodate. The aldehyde is then converted to the
 corresponding acid as described in Reaction Scheme D. The olefin can
 be cleaved to the carboxylic acid directly with ruthenium tetroxide and
 periodate.

REACTION SCHEME F



As depicted in Reaction Scheme F, esters can be prepared by

5 reaction of an acid chloride with the hydroxy group in a basic solvent such as pyridine. It should be understood that R^4 is used to represent a portion of the R^4 definition. The acid chlorides, when not purchased, are prepared by stirring the carboxylic acids in reagents such as oxalyl chloride or thionyl chloride. Esters may also be prepared by reaction of

10 the acid chloride and the hydroxy group with silver cyanide (AgCN) in an aprotic solvent such as HMPA. C1 sulfonate derivatives are prepared in a similar manner by reaction with sulfonyl chlorides.


Carbonate and carbamate derivatives are prepared by first reacting the alcohol derivative with carbonyldiimidazole (CDI) to obtain

15 the imidazolecarbonyl intermediate which is then reacted with an alcohol (R^4OH) or amine ($\text{R}^4\text{R}^4\text{NH}$) to give the corresponding carbonate or carbamate derivatives. In an alternate approach, reaction

of the hydroxy group with 4-nitrochloroformate provides the 4-nitrophenylcarbonate which then can be reacted with amines to give carbamates or with alcohols to give carbonate derivatives. Carbamate derivatives are also prepared with commercially available carbamoyl chlorides or isocyanates.

Ether derivatives can also be prepared. A particularly useful procedure involves reacting an alcohol with trifluoromethane sulfonic anhydride (Tf_2O , triflic anhydride) in dichloromethane at reduced temperature, preferably -78°C to obtain the preformed triflate. To this solution is added the alcohol, the reaction mixture is warmed to room temperature and stirring is continued until reaction is complete. Ethers may also be prepared by heating a mixture of alcohol, the appropriate alkylhalide and an excess of silver oxide (Ag_2O) in an aprotic inert solvent such as THF.

REACTION SCHEME G




In Reaction Scheme G, the hydroxy group is converted to the corresponding amino group by well-known procedures. The hydroxy group is first derivatized as the methanesulfonate with methanesulfonyl chloride in pyridine or in THF or dichloromethane with triethylamine or pyridine. This intermediate is then reacted with sodium azide in a polar aprotic solvent such as dimethylformamide. Alternatively, lithium azide in THF can be used. The azide displacement occurs with inversion, so that the beta hydroxy group gives the alpha azide derivative

and vice versa. Finally, the azide is reduced to the corresponding amino group by hydrogenation using a palladium catalyst. This and other procedures are described in March, J.; *Advanced Organic Chemistry*, 4th ed., John Wiley & Sons, New York, pp. 428 and 1219 (1992).

5

REACTION SCHEME H



As depicted in Reaction Scheme H, amides are prepared by
 10 reaction of an acid chloride with the amino group in a basic solvent such
 as pyridine. The acid chlorides, when not purchased, are prepared by
 stirring the carboxylic acids in reagents such as oxalyl chloride or
 thionyl chloride. Amides may also be prepared from carboxylic acids by
 using coupling reagents such as dicyclohexyl carbodiimide as reviewed
 15 by Bodanski, *The Practice of Peptide Synthesis*, Springer, New York,
 (1984). C1 sulfonamide derivatives are prepared in a similar manner by
 reaction with sulfonyl chlorides.

Urea derivatives are prepared by first reacting the C4 amino derivative with carbonyldiimidazole (CDI) to obtain the imidazolecarbonyl intermediate which is then reacted with an amine ($R^4'R^4''NH$) to give the corresponding urea derivatives. In an alternate approach, reaction of the amino group with 4-nitrochloroformate provides the 4-nitrophenyl carbamate which then can be reacted with amines to give urea derivatives. Urea derivatives are also prepared with commercially available carbamoyl chlorides or isocyanates.

Carbamate derivatives are prepared in a similar manner. Reacting the C4 amino derivative with carbonyldiimidazole (CDI) gives the imidazolecarbonyl intermediate which is then reacted with an alcohol to give the corresponding carbamate derivatives. In an alternate approach, reaction of the amino group with 4-nitrochloroformate provides the 4-nitrophenylcarbamate which then can be reacted with alcohols. Carbamate derivatives are also prepared with chloroformates. For instance, reaction with ethylchloroformate will give the ethylcarbamate derivative.

The amino group can be alkylated by reductive amination procedures. For instance, the amino group is reacted with an aldehyde or ketone in the presence of sodium cyanoborohydride or sodium triacetoxyborohydride. This transformation may also be accomplished with hydrogen and a catalyst.

UTILITY

The present invention is related to compounds of Formula I, including but not limited to those specified in the examples, which are useful in a mammalian subject for the treatment and prevention of immunemediated diseases such as the resistance by transplantation of organs or tissue such as heart, kidney, liver, medulla ossium, skin, cornea, lung, pancreas, intestinum tenue, limb, muscle, nervous, duodenum, small-bowel, pancreatic-islet-cell, including xeno

transplants, etc.; graft-versus-host diseases brought about by medulla
ossium transplantation; autoimmune diseases such as rheumatoid
arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis,
multiple sclerosis, myasthenia gravis, type I diabetes uveitis, juvenile-
5 onset or recent-onset diabetes mellitus, posterior uveitis, allergic
encephalomyelitis, glomerulonephritis, and the like; and further
infectious diseases caused by pathogenic microorganisms. Further uses
may include the treatment and prophylaxis of inflammatory and
hyperproliferative skin diseases and cutaneous manifestations of
10 immunologically mediated illnesses, such as psoriasis, atopic
dermatitis, contact dermatitis and further eczematous dermatitises and
further eczematous dermatitises, seborrhoeis dermatitis, Lichen
planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa,
urticaria, angioedemas, vasculitides, erythemas, cutaneous
15 eosinophilias, Lupus erythematosus, acne and Alopecia areata; various
eye diseases (autoimmune and otherwise) such as keratoconjunctivitis,
vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis,
herpetic keratitis, conical cornea, dystrophia epithelialis corneae,
corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves'
20 ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, etc.;
pollen allergies, reversible obstructive airway disease, which includes
condition such as asthma (for example, bronchial asthma, allergic
asthma, intrinsic asthma, extrinsic asthma and dust asthma),
particularly chronic or inveterate asthma (for example, late asthma and
25 airway hyper-responsiveness), bronchitis and the like; inflammation of
mucous and blood vessels such as gastric ulcers, vascular damage
caused by ischemic diseases and thrombosis, ischemic bowel diseases,
inflammatory bowel diseases, necrotizing enterocolitis, intestinal
lesions associated with thermal burns and leukotriene B₄-mediated
30 diseases; intestinal inflammations/allergies such as Coeliac diseases,
proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease and

- ulcerative colitis; food-related allergic diseases which have symptomatic manifestation remote from the gastrointestinal tract (e.g. migraine, rhinitis and eczema); renal diseases such as interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome and diabetic
- 5 nephropathy; nervous diseases such as multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis, mononeuritis and radiculopathy; endocrine diseases such as hyperthyroidism and Basedow's disease; hematic diseases such as pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic
- 10 thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia and anerythroplasia; bone diseases such as osteoporosis; respiratory diseases such as sarcoidosis, fibroid lung and idiopathic interstitial pneumonia; skin disease such as dermatomyositis, leukoderma
- 15 vulgaris, ichthyosis vulgaris, photoallergic sensitivity and cutaneous T cell lymphoma; circulatory diseases such as arteriosclerosis, atherosclerosis, aortitis syndrome, polyarteritis nodosa and myocardosis; collagen diseases such as scleroderma, Wegener's granuloma and Sjogren's syndrome; adiposis; eosinophilic fascitis;
- 20 periodontal disease such as lesions of gingiva, periodontium, alveolar bone and substantia ossea dentis; nephrotic syndrome such as glomerulonephritis; male pattern alopecia or alopecia senilis by preventing epilation or providing hair germination and/or promoting hair generation and hair growth; muscular dystrophy; Pyoderma and
- 25 Sezary's syndrome; Addison's disease; active oxygen-mediated diseases, as for example organ injury such as ischemia-reperfusion injury of organs (such as heart, liver, kidney and digestive tract) which occurs upon preservation, transplantation or ischemic disease (for example, thrombosis and cardiac infraction); intestinal diseases such as
- 30 endotoxin-shock, pseudomembranous colitis and colitis caused by drug or radiation; renal diseases such as ischemic acute renal insufficiency

and chronic renal insufficiency; pulmonary diseases such as toxinosis caused by lung-oxygen or drug (for example, paracort and bleomycins), lung cancer and pulmonary emphysema; ocular diseases such as cataracta, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring and corneal alkali burn; dermatitis such as erythema multiforme, linear IgA ballous dermatitis and cement dermatitis; and others such as gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution (for example, air pollution), aging, carcinogenis, metastasis of carcinoma and hypobaropathy; disease caused by histamine or leukotriene-C4 release; Behcet's disease such as intestinal-, vasculo- or neuro-Behcet's disease, and also Behcet's which affects the oral cavity, skin, eye, vulva, articulation, epididymis, lung, kidney and so on. Furthermore, the compounds of the invention are useful for the treatment and prevention of hepatic disease such as immunogenic diseases (for example, chronic autoimmune liver diseases such as the group consisting of autoimmune hepatitis, primary biliary cirrhosis and sclerosing cholangitis), partial liver resection, acute liver necrosis (e.g. necrosis caused by toxin, viral hepatitis, shock, or anoxia), B-virus hepatitis, non-A/non-B hepatitis, cirrhosis (such as alcoholic cirrhosis) and hepatic failure such as fulminant hepatic failure, late-onset hepatic failure and "acute-on-chronic" liver failure (acute liver failure on chronic liver diseases), and moreover are useful for various diseases such as augmentation of chemotherapeutic effect, preventing or treating activity of cytomegalovirus infection, particularly HCMV infection, and antiinflammatory activity.

The compounds of the present invention may also be used in the treatment of immunodepression or a disorder involving immunodepression, such as AIDS, cancer, senile dementia, trauma (including wound healing, surgery and shock) chronic bacterial infection, and certain central nervous system disorders.

Using the methodologies described below, representative compounds of the invention were evaluated and found to exhibit activity thereby demonstrating and confirming the utility of the compounds of the invention as $K_v1.3/K_v1.5$ inhibitors and therefore as

5 immunosuppressants and antiarrhythmics.

ASSAYS

T CELL IL-2 ASSAY

Peripheral blood mononuclear (MNC) cells from healthy

10 donors were separated by density centrifugation with ficoll-hypaque (LSM, Organon Teknika, Durham, NC), followed by rosetted with neuraminidase treated sheep red blood cells (SRBC). After another centrifugation with leucocyte separation medium (LSM), the SRBC of the rosetted T cells were then lysed with ammonium chloride lysing buffer

15 (GIBCO, Grand Island, NY). Such purified T cells were resuspended at 3×10^6 /mL in RPMI 1640 culture medium (GIBCO) supplemented with 10% fetal calf serum (Sigma, St. Louis, MO), 100 mM glutamine, 1 mM sodium pyruvate, 0.1 mM non-essential amino acids, and 1 % penn-strep (GIBCO). The cell suspension was immediately distributed into 96

20 well round-bottom microculture plates (Costar) at 200 μ L/well. The various dilutions of test compound were then added in triplicate wells at 25 μ L/well, incubated for 30 min at 37°C. Ionomycin (125 ng/mL), and PMA (1 or 5 ng/mL), were added to the appropriate wells. The culture plates were then incubated at 37°C in a humidified atmosphere of 5%

25 CO₂ - 95% air for 18-24 hours. The supernatants were removed, and assayed for IL-2 with an IL-2 capture ELISA, using monoclonal anti-IL-2, and biotinylated goat anti-IL-2 antibodies (unconjugated antibodies purchased from R&D System, Minneapolis, MN). The ELISA was developed with streptavidin conjugated peroxidase (Zymed, San

30 Francisco, CA) and substrate for peroxidase (Sigma). Mean OD and units of IL-2 of the replicate wells were calculated from standard curve,

created with recombinant IL-2 (Collaborative Biomedical Products, Bedford, MA) and the results were expressed as concentration of compound required to inhibit IL-2 production of T cells by 50%.

5

T CELL PROLIFERATION ASSAY

Peripheral blood mononuclear cells (MNC) from healthy donors were separated by density centrifugation with ficoll-hypaque (LSM, Organon Teknika, Durham, NC). After washing the MNC with complete media (RPMI 1640 medium with 5% fetal calf serum, 100 mM glutamine, 1 mM sodium pyruvate, 0.1 mM non-essential amino acid, and 1% penn-strep, obtained from GIBCO, Grand Island, NY), they were then irradiated at 7500 RADS, and resuspended at $4-4.5 \times 10^6$ cells/mL in complete media. Another aliquot of MNC were rosetted with neuraminidase treated SRBC. After another centrifugation with LSM, the sheep red blood cells (SRBC) of these rosetted T cells were then lysed with ammonium chloride lysing buffer (GIBCO, Grand Island, NY). After washing 2X with complete media, these purified T cells were also resuspended at $2-2.5 \times 10^6$ cells/mL in complete media. The various dilutions of the compound were added in triplicates at 50 μ l/well of a 96 well flat-bottom microculture plate (Costar, Cambridge, MA). T cell suspension was then immediately distributed into the wells at 100 μ l/well. After incubating the cells with compound for 30 min. at 37°C in a humidified atmosphere of 5% CO₂ - 95% air, 20 μ l/well of anti-CD3 (Ortho Diagnostic, NJ) at final conc. of 0.3 ng/mL was added, followed by 50 μ l of the irradiated MNC. The culture plates were then incubated at 37°C in a humidified atmosphere of 5% CO₂ - 95% air for 72 hours. The proliferation of T lymphocytes was assessed by measurement of tritiated thymidine incorporation. During the last 18-24 hrs. of culturing, the cells were pulse-labeled with 2 μ Ci/well of tritiated thymidine (NEN, Cambridge, MA). The cultures were harvested on glass fiber filters

30

using a multiple sample harvester (MACH-II, Wallac, Gaithersburg, MD). Radioactivity of filter discs corresponding to individual wells was measured by standard liquid scintillation counting methods (Betaplate Scint Counter, Wallac). Mean counts per minute of replicate wells were
5 calculated and the results were expressed as concentration of compound required to inhibit tritiated thymidine uptake of T cells by 50%.

K_v1.3-RUBIDIUM EFFLUX ASSAY

CHO cells transfected with K_v1.3 channels at site densities
10 of approximately 40,000 sites/cell are plated into 96 well culture plates and maintained in Iscove's Modified Dulbecco's Medium (IMDM, with L-Glutamine and HEPES, JRH Biosciences). Cells are incubated overnight with ⁸⁶Rb⁺ (3 μCi/mL, Dupont-NEN) in the glutamine supplemented IMDM. After aspiration of the media, 100 μL of Low K
15 Buffer (in mM, 6.5 KCl, 125 NaCl, 1 CaCl₂, 2 MgCl₂, 10 HEPES, pH adjusted to 7.2 with NaOH) is added to each well followed by 100 μL test samples in Low K Buffer also containing 0.2% BSA and 2 mM ouabain. Samples are tested at either 1 μg/mL for routine screening or at a variety of concentrations encompassing at least 1/10 to 10 times the putative
20 IC₅₀ of test compound to determine potency. After a fixed preincubation time, which is usually 10 min, the samples are aspirated. The K_v1.3 channels are opened by depolarization of the cells with High K Buffer (final concentrations, in mM, 63.25 KCl, 68.25 NaCl, 1 CaCl₂, 2 MgCl₂, 10 HEPES, pH adjusted to 7.2 with NaOH) also containing test
25 compounds. To measure ⁸⁶Rb⁺ efflux through the channels, aliquots of 100 μL are taken from each well after a given time and added to plates containing 100 μL MicroScint-40 (Packard) for counting by liquid scintillation techniques. MicroScint-40 (100 μL) is then added to each well of the cell plate to determine the remaining ⁸⁶Rb⁺ activity. The
30 efflux counts are normalized for the total amount of ⁸⁶Rb⁺ that was in

the cells by adding the efflux counts to the cell plate counts. Activity is determined by % inhibition of the efflux window that is established using a saturating concentration of margatoxin (MgTX), a 39 amino acid peptide that is a potent blocker of $K_v1.3$ channels ($IC_{50} = 100$ pM).

5

$K_v1.5$ -RUBIDIUM EFFLUX ASSAY

The $K_v1.5$ $^{86}Rb^+$ (a potassium ion surrogate) efflux assay utilizes HEK 293 cells engineered to stably express the human $K_v1.5$ potassium channel. Cells are seeded at a density of 25000 cells/well in poly-D-lysine coated 96-well Packard CulturPlates one to three days prior to the assay and loaded with $^{86}Rb^+$ (0.05 microcuries/well) the day before assay. On the day of assay, plates are washed three times using a Skatron 96-well plate washer and two hundred microliters of low KCl buffer (125mM NaCl, 6.5mM KCl, 1mM $CaCl_2$, 2mM $MgCl_2$, 0.02% bovine serum albumin, 10mM HEPES, pH 7.2) with or without inhibitor is added to each well. After ten minutes at room temperature, the plates are washed one time with low KCl buffer and two hundred microliters of high KCl buffer (same as low KCl buffer except KCl is 60mM and NaCl is 65mM) with or without inhibitor is added to the appropriate wells to activate the $K_v1.5$ channel. The plates are incubated for an additional ten minutes at room temperature at which time one hundred microliters of each supernatant is transferred to a 96-well Packard OptiPlate. The cell plates are immediately washed one time with low KCl buffer followed by the addition of one hundred microliters of low KCl buffer to each well. One hundred microliters of Microscint scintillation cocktail is added to each well of the supernatant and cell plates and radioactivity is determined on a Packard TopCount scintillation counter. The reduction of supernatant $^{86}Rb^+$ is used to quantitate the degree of inhibition of the $K_v1.5$ potassium channel.

30

Di-TRITIUM CORREOLIDE (DiTC) BINDING ASSAY

Activity of compounds can be evaluated by determining their effect on DiTC binding to either Kv1.3 or Kv1.5 channels expressed in human embryonic kidney cells (HEK). The Kv1.3 and Kv1.5 channels used are
5 cloned human channels. DiTC is also known as L-765,910 and is a natural product analogue which binds specifically to Kv1.x channels, such as Kv1.3 and Kv1.5. In general, compounds are incubated at given concentrations in the presence of 20 nM DiTC (10 μ M cold ligand is used to define non-specific binding) and either HEK/Kv1.3 or HEK/Kv1.5
10 membranes, in a buffer containing 135 mM NaCl, 4.6 mM KCl, 20 mM Tris, pH = 7.4 (tris[hydroxymethyl]aminomethane), and 0.02% bovine serum albumin (BSA). Binding is allowed to reach equilibrium by incubation of the samples at room temperature for 24 hrs. Separation of bound from free ligand is achieved by filtration through GF/C filters and
15 washing with ice-cold buffer containing 100 mM NaCl, 20 mM Tris (pH = 7.4), and 0.04% BSA. Scintillation fluid is added to the samples and radioactivity associated with filters determined by liquid scintillation techniques. Specific DiTC binding is the difference between total and non-specific binding. The activity of a given compound is assessed
20 relative to an untreated control.

HEK cells expressing either human Kv1.3 or Kv1.5 channels were grown by Analytical Biological Services in bioreactors containing MEM supplemented with FBS, Penicillin, Streptomycin, and Geneticin. Seven tubes of frozen cell pellets (25 L of cells) were then thawed and 20 mL
25 Homogenization Buffer was added to each tube. The contents of the tubes were pooled into a 50 mL glass/Teflon homogenizer. The cells were homogenized for 10 strokes (500 rpm) and transferred to 50 mL tubes. The tubes were then centrifuged at 1000 rpm for 5 min at 4 °C (253 xg, Beckman GPR). The supernatant was collected and set aside on ice.
30 The pellets were resuspended in a total of 40 mL Lysis Buffer, homogenized as described above, and the homogenate was centrifuged

as described above. The supernatant was added to the set aside supernatant. The pooled supernatant was then centrifuged at 40,000 rpm for 45 min at 4 °C (186,000 xg, Beckman 45TI). The pellet was resuspended in 70 mL Storage Buffer by homogenization as described
5 above. Aliquots were flash frozen using liquid nitrogen and stored at -70 °C. (Homogenization Buffer: 250 mM Sucrose, 5 mM MgCl₂, 20 mM Tris, pH = 7.4; Storage Buffer: 100 mM NaCl, 20 mM HEPES, pH = 7.4).

DOSAGE FORMS

10 As an immunosuppressive, these compounds are useful in the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and/or related afflictions, diseases and illnesses.

The compounds of this invention can be administered for
15 the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and/or related afflictions, diseases and illnesses according to the invention by any means that effects contact of the active ingredient compound with the site of action in the body of a warm-blooded animal. For example, administration, can be oral,
20 topical, including transdermal, ocular, buccal, intranasal, inhalation, intravaginal, rectal, intracisternal and parenteral. The term "parenteral" as used herein refers to modes of administration which include subcutaneous, intravenous, intramuscular, intraarticular injection or infusion, intrasternal and intraperitoneal.

25 The compounds can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but are generally administered with a pharmaceutical carrier selected on the basis of the chosen route of
30 administration and standard pharmaceutical practice.

For the purpose of this disclosure, a warm-blooded animal is a member of the animal kingdom possessed of a homeostatic mechanism and includes mammals and birds.

The dosage administered will be dependent on the age,
5 health and weight of the recipient, the extent of disease, kind of concurrent treatment, if any, frequency of treatment and the nature of the effect desired. Usually, a daily dosage of active ingredient compound will be from about 1-500 milligrams per day. Ordinarily, from 10 to 100 milligrams per day in one or more applications is effective to obtain
10 desired results. These dosages are the effective amounts for the treatment of autoimmune diseases, the prevention of rejection of foreign organ transplants and/or related afflictions, diseases and illnesses.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, troches, dragées, granules and
15 powders, or in liquid dosage forms, such as elixirs, syrups, emulsions, dispersions, and suspensions. The active ingredient can also be administered parenterally, in sterile liquid dosage forms, such as dispersions, suspensions or solutions. Other dosage forms that can also be used to administer the active ingredient as an ointment, cream,
20 drops, transdermal patch or powder for topical administration, as an ophthalmic solution or suspension formation, i.e., eye drops, for ocular administration, as an aerosol spray or powder composition for inhalation or intranasal administration, or as a cream, ointment, spray or suppository for rectal or vaginal administration.

25 Gelatin capsules contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous
30 release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the

tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

5 In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents,
10 and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or
15 propylparaben, and chlorobutanol.

Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field.

For administration by inhalation, the compounds of the
20 present invention may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The compounds may also be delivered as powders which may be formulated and the powder composition may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery system for
25 inhalation is a metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of a compound of Formula I in suitable propellants, such as fluorocarbons or hydrocarbons.

For ocular administration, an ophthalmic preparation may be formulated with an appropriate weight percent solution or
30 suspension of the compounds of Formula I in an appropriate ophthalmic vehicle, such that the compound is maintained in contact

with the ocular surface for a sufficient time period to allow the compound to penetrate the corneal and internal regions of the eye.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as follows:

5

CAPSULES

A large number of unit capsules are prepared by filling standard two-piece hard gelatin capsules each with 100 milligrams of powdered active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.

10

SOFT GELATIN CAPSULES

A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules are washed and dried.

15

TABLETS

A large number of tablets are prepared by conventional procedures so that the dosage unit is 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275 milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

20

25

INJECTABLE

A parenteral composition suitable for administration by injection is prepared by stirring 1.5% by weight of active ingredient in

30

10% by volume propylene glycol. The solution is made to volume with water for injection and sterilized.

SUSPENSION


5 An aqueous suspension is prepared for oral administration so that each 5 milliliters contain 100 milligrams of finely divided active ingredient, 100 milligrams of sodium carboxymethyl cellulose, 5 milligrams of sodium benzoate, 1.0 grams of sorbitol solution, U.S.P., and 0.025 milliliters of vanillin.

10 The same dosage forms can generally be used when the compounds of this invention are administered stepwise or in conjunction with another therapeutic agent. When drugs are administered in physical combination, the dosage form and administration route should be selected depending on the compatibility
15 of the combined drugs. Thus the term coadministration is understood to include the administration of the two agents concomitantly or sequentially, or alternatively as a fixed dose combination of the two active components.

 The following examples illustrate the preparation of the
20 compounds of Formula I and as such are not to be considered as limiting the invention set forth in the claims appended hereto.

EXAMPLE 1


25 1-(2-methoxyphenyl)-1-oxo-2-aza-4-phenylbutane



To a solution of 0.5 gm of phenethylamine (4.12 mmol) in 5 mL of dichloromethane and 0.978 mL of pyridine (8.2 mmol) at 0°C was added 1.39 gm of 2-methoxybenzoyl chloride (1.14 mL, 8.2 mmol). The reaction was allowed to warm to room temperature and was stirred for 4
5 hr. The reaction mixture was concentrated, 5mL of heptane were added and the mixture was evaporated to dryness this being repeated one more time. The residue was purified by HPLC (RCM SepPak, silica 25x100, 3.6% CH₃CN, 14.4% tBuOCH₃, 82% hexane) to give the title compound.
¹H NMR (CDCl₃) δ 2.95 (m, 2H), 3.8 (s,m, 5H), 6.95 (d, 1H), 7.1 (t, 1H), 7.4
10 (m, 6H), 7.9 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 256 (M⁺1).

EXAMPLE 2

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-phenylbutane



15

Step 1 3,3-diethyl-3-phenethylamine

To a solution of 0.25 gm of 2,2-diethylphenylacetonitrile (1.44
20 mmol, purchased from Sigma-Aldrich Rare Chemicals) in 10 mL of THF at room temperature, was added 7.22 mL of a 1 M solution of lithium aluminum hydride (LAH) in THF (7.22 mmol). After stirring for 12 hr, the reaction was quenched with with the successive addition of 1 mL of water, 1 mL of 15% NaOH solution and a 1 mL of water. The
25 aluminum salts were removed by filtration and washed with 100mL of ethylacetate. The filtrate was dried over MgSO₄, filtered and

concentrated. The title compound was used without further purification.


5 Step 2 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-phenylbutane

To a solution of 3,3-diethyl-3-phenethylamine (Step 1) in 5 mL of dichloromethane at room temperature was added 0.405 mL of 2-methoxybenzoyl chloride (2.7 mmol), 0.164 mL of pyridine (2.02 mmol) and 10 mg of 4,4-dimethylaminopyridine (DMAP). After stirring for 12
10 hr, the reaction mixture was concentrated in vacuo and purified by chromatography (silica, 2:1 hexanes: ethyl acetate). The residue was repurified by HPLC (RCM SepPak, silica 25x100, 2.6% CH₃CN, 10.6% tBuOCH₃, 86.8% hexane) to afford the title compound.

¹H NMR (CDCl₃) δ 0.8 (t, 6H), 1.8 (m, 4H), 1.8 (m, 1H), 3.55 (s, 3H), 3.8 (d, 2H), 6.85 (d, 1H), 7.05 (t, 1H), 7.35 (m, 1H), 7.4 (m, 5H), 7.65 (br s, 1H), 8.2
15 (dd, 1H). Mass Spectrum m/e 312 (M⁺).

EXAMPLE 3

20 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-dimethyl-4-phenylbutane



Step 1 2,2-dimethylphenylacetonitrile

To a solution of 1 gm of phenylacetonitrile (8.5 mmol) in 10
25 mL of anhydrous DMSO, at 0 °C, was added 0.043 gm of NaH (17.1 mmol). After stirring for 0.5 hr, 1.2 mL of methyl iodide (19.6 mmol) was slowly added to the reaction mixture. After 6 hours, TLC indicated that the reaction was complete. The reaction was quenched with H₂O.

The aqueous layer was extracted 3 times with ether. The combined organic fractions were dried over MgSO_4 , filtered through a thin layer of silica gel and concentrated *in vacuo* to provide the title compound which was used without further purification.


- 5 2,2-dimethylphenylacetonitrile was converted to the title compound according to procedures described in Example 2.

^1H NMR (CDCl_3) δ 1.4 (s, 6H), 3.55 (s, 3H), 3.75 (d, 2H), 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 1H), 7.4 (m, 5H), 7.65 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 284 (M^+1).

10

EXAMPLE 4

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-dibenzyl-4-phenylbutane




- 15 The title compound was prepared from 2,2-dibenzyl acetonitrile (Sigma-Aldrich Rare Chemicals) according to procedures described in Example 2.

^1H NMR (CDCl_3) δ 3.15 (s, 4H), 3.55 (s, 3H), 3.8 (d, 2H), 6.9 (m, 5H), 7.1 (m, 1H), 7.15 (m, 5H), 7.35 (m, 5H), 7.4 (m, 1H), 8.0 (br s, 1H), 8.2 (dd, 1H).

- 20 Mass Spectrum m/e 436 (M^+1).

EXAMPLE 5

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diallyl-4-phenylbutane



Step 1 2,2-diallylphenylacetonitrile

The title compound was prepared from phenylacetonitrile,
5 allyl chloride and NaH as described in Example 3.

Step 2 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diallyl-4-phenylbutane


The title compound was prepared from 2,2-diallylphenyl
10 acetonitrile as described in Example 2.

^1H NMR (CDCl_3) δ 0.8 2.55 (m, 4H), 3.6 (s, 3H), 3.8 (d, 2H), 5.1 (m, 4H),
5.7 (m, 2H), 6.85 (d, 1H), 7.05 (t, 1H), 7.30 (m, 1H), 7.4 (m, 5H), 7. (br 70s,
1H), 8.2 (dd, 1H). Mass Spectrum m/e 336 (M^+1).

15

EXAMPLE 6

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-di-n-propyl-4-phenylbutane



20


A mixture of 0.2 gm of 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diallyl-4-phenylbutane (Example 5) in 10 mL of methanol and 0.025 gm of 5% Pd/C was hydrogenated at 40 psi (Parr Apparatus). After 12 hr, the

reaction mixture was filtered through a thin pad of silica and concentrated in vacuo to afford the title compound.

^1H NMR (CDCl_3) δ 0.85 (t, 6H), 1.15 (m, 2H), 1.3 (m, 2H), 1.7 (m, 4H), 3.55 (s, 3H), 3.8 (d, 2H), 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 1H), 7.4 (m, 5H), 7.65 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 340 (M^+).

EXAMPLE 7

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-di-(3-hydroxypropyl)-4-phenylbutane



10


To 0.512 gm of 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diallyl-4-phenylbutane (Example 5, 1.5 mmol) in a nitrogen atmosphere, was added 12 mL of a 0.5 M solution of 9-borabicyclo(3.3.1)nonane (9-BBN) in hexanes, and the reaction mixture was stirred at 35 °C. After 4 hr, the reaction mixture was cooled to 0 °C and to it was added 5 mL of 15% NaOH solution, 2 mL of hydrogen peroxide solution (30% in water) and 3 mL of THF. After 3 hr at room temperature, the reaction mixture was extracted 3 times with ethyl acetate. The combined organic fractions were washed with 1N HCl and sat. NaHCO_3 , dried over MgSO_4 , filtered and concentrated. The residue was purified by chromatography (silica, 4:1, ethyl acetate: hexanes) to give the title compound.

^1H NMR (CDCl_3) δ 1.4 (m, 2H), 1.6 (m, 2H), 1.8 (m, 4H), 2.2 (br s, 2H), 3.55 (s, 3H), 3.6 (m, 4H), 3.8 (d, 2H), 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 1H), 7.4 (m, 5H), 7.65 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 372 (M^+).

25

EXAMPLE 8

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-di-(3-acetoxypentyl)-4-phenylbutane



5

A solution of 0.085 gm of 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-di-(3-hydroxypentyl)-4-phenylbutane (Example 7, 0.23 mmol), 0.082 mL of acetic anhydride (0.69 mmol) and 0.056 mL of pyridine (0.69 mmol) in 2 mL of DMF was stirred for 12h. The reaction mixture was concentrated, 5mL of heptane were added and the mixture was evaporated to dryness this being repeated one more time. The residue was purified by HPLC (RCM SepPak, silica 25x100, 5% CH₃CN, 20% tBuOCH₃, 75% hexane) to give the title compound.


¹H NMR (CDCl₃) δ 1.5 (m, 2H), 1.6 (m, 2H), 1.8 (m, 4H), 2.0 (s, 6H), 3.55 (s, 3H), 3.8 (d, 2H), 4.0 (t, 4H) 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 1H), 7.4 (m, 5H), 7.65 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 456 (M⁺1).

15

EXAMPLE 9

20

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-di-(2-hydroxyethyl)-4-phenylbutane



A solution of 500 mg of 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diallyl-4-phenylbutane (1.5 mmol) in 5 mL of methanol and 5mL of methylene chloride was cooled to -78 °C. Ozone was bubbled through the reaction mixture for 20 min. until tlc (silica, 9:1 CH₂Cl₂:CH₃OH)


- 5 showed no more starting material. 0.5 gm of NaBH₄ was then added to the reaction mixture and it was warmed to room temperature. After 0.5 hr 20 mL of 1N aqueous HCl were carefully added and the mixture extracted with CH₂Cl₂. The combined organic extracts were washed with brine, dried over magnesium sulfate, filtered, and concentrated.
- 10 The residue was purified by chromatography (silica, 9:1 ethyl acetate : hexanes) to give the title compound.

¹H NMR (CDCl₃) δ 2.0 (t, 4H), 3.55 (m, 4H), 3.7 (s, 3H), 3.9 (s, 2H), 4.0 (t, 4H), 7.0 (m, 2H), 7.3 (m, 1H), 7.4 (m, 6H), 8.0 (dd, 1H). Mass Spectrum m/e 344 (M⁺1).

15

EXAMPLE 10

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-di-(2-acetoxyethyl)-4-phenylbutane




20

The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-di-(2-hydroxyethyl)-4-phenylbutane (Example 9) according to procedures described in Example 8.

- ¹H NMR (CDCl₃) δ 2.0 (s, 6H), 2.2 (t, 4H), 3.6 (s, 3H), 3.9 (d, 2H), 4.1 (m, 4H) 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 1H), 7.4 (m, 5H), 7.65 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 428 (M⁺1).
- 25

EXAMPLE 11

1-(2-methoxyphenyl)-1-oxo-2-aza-(R)-ethyl-4-phenylbutane



5

Step 1 (R)-(-)-2-phenylbutanol

To a suspension of 2.3 gm of LAH (60.6 mmol) in 70 mL of THF was slowly added a solution of 5 gm of (R)-(-)-2-phenylbutyric acid
10 (30.45 mmol, Aldrich) in 30 mL of THF. The reaction mixture was heated to 60 °C for 3 hr. The reaction was cooled to 0 °C and to it was successively added 2 mL of water, 2 mL of 1N KOH and 2mL of water. The aluminum salts were removed by filtration and washed with 100mL of ethylacetate. The filtrate was dried over MgSO₄, filtered and
15 concentrated to give the title compound which was used without purification.

¹H NMR (CDCl₃) δ 0.9 (t, 3H), 1.6 (m, 2H), 1.8 (m, 2H), 2.2 (bs, 1H), 2.7 (m, 1H), 3.75 (m, 2H), 7.3 (m, 5H).

20 Step 2 (R)-(-)-2-phenylbutyl amine

To a solution of 4.7 gm of (R)-(-)-2-phenylbutanol (31.78 mmol) and 7.6 mL of pyridine (95 mmol) in 40 mL of CH₂Cl₂ was added 10 gm of methanesulfonic anhydride (64 mmol). After stirring at room
25 temperature for 5 hr, 100 mL of water was added and the mixture was extracted 3 times with ether. The combined organic fractions were washed with 1N HCl, sat. NaHCO₃ and sat. NaCl solutions. The


organic fractions were dried over MgSO_4 , filtered through a thin layer of silica and concentrated in vacuo. The methanesulfonate was used without further purification. ^1H NMR (CDCl_3) δ 0.85 (t, 3H), 1.65 (m, 2H), 1.85 (m, 2H), 2.75 (s, 3H), 2.9 (m, 1H), 4.35 (d, 2H), 7.3 (m, 5H). A solution of the methanesulfonate, 10 gm of sodium azide (154 mmol) and a trace amount of sodium iodide in 50 mL of DMF was stirred at 80 °C for 12 hr. The reaction mixture was then cooled and 200 mL of ether was added. The mixture was washed 3 times with water, then with sat. NaCl solution, dried over MgSO_4 , filtered and concentrated. To a solution of the azide in 20 mL of THF was added 50 mL of a 1M solution of LAH in THF and the reaction mixture was stirred at room temperature. After 2 hours, the reaction was quenched with successive additions of 5 mL of water, 5 mL of 10% KOH and 5 mL of water. The aluminum salts were removed by filtration and washed with 200 mL of ethylacetate. The filtrate was dried over MgSO_4 , filtered and concentrated. The residue was purified by vacuum distillation in a Kugelrohr apparatus (bp 145 °C, 0.2 Torr) to give the title compound as a clear liquid.

20 Step 3 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-ethyl-4-phenylbutane
The title compound was prepared from 0.2 gm of (R)-(-)-2-phenylbutyl amine and 2-mL of 4 methoxybenzoyl chloride according to procedures described in Example 2.

^1H NMR (CDCl_3) δ 0.87 (t, 3H), 1.66 (m, 1H), 1.8 (m, 1H), 2.8 (m, 1H), 3.4 (m, 1H), 3.55 (s, 3H), 4.05 (m, 1H), 6.87 (d, 1H), 7.03 (t, 1H), 7.3 (m, 6H), 7.75 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 284 (M^+), α_D (CHCl_3) = +30.2.

EXAMPLE 12

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-ethyl-4-phenylbutane




5 The title compound was prepared from (S)-(+)-2-phenyl butyric acid according to procedures described in Example 11.

^1H NMR (CDCl_3) same as Example 11, Step 3 Mass Spectrum m/e 284 (M^+1), α_D (CHCl_3) = -29.7.

10

EXAMPLE 13

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-methyl-4-phenylbutane




15 The title compound was prepared from (R)-(-)-2-phenyl propionic acid according to procedures described in Example 11.

^1H NMR (CDCl_3) δ 1.4 (d, 3H), 3.1 (m, 1H), 3.5 (m, 1H), 3.6 (s, 3H), 3.95 (m, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.35 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H).
Mass Spectrum m/e 270 (M^+1), α_D (CHCl_3) = +34.25.

20

EXAMPLE 14

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-methyl-4-phenylbutane




The title compound was prepared from (S)-(+)-2-phenyl propionic acid according to procedures described in Example 11.

^1H NMR (CDCl_3) same as Example 13. Mass Spectrum m/e 270 (M^+),

5 α_D (CHCl_3) = -35.05.

EXAMPLE 15

1-(2-methoxyphenyl)-1-oxo-2-aza--(S)-4-i-propyl-4-phenylbutane




10

The title compound was prepared from (S)-(+)-2-phenyl-3-methylbutyric acid (Aldrich) according to procedures described in Example 11.

^1H NMR (CDCl_3) δ 0.75 (d, 3H), 1.0 (d, 3H), 1.85 (m, 1H), 2.5 (m, 1H), 3.35
 15 (m, 4H), 4.2 (m, 1H), 6.7 (d, 1H), 6.95 (t, 1H), 7.2 (m, 6H), 7.6 (br s, 1H),
 8.14 (dd, 1H). Mass Spectrum m/e 298 (M^+), α_D (CHCl_3) = -19.8.

EXAMPLE 16

20 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-allyl-4-phenylbutane



5 Step 1 3-Phenylacetyl-4-(S)-benzyl-2-oxazolidinone

A solution of 10 gm of phenylacetic acid (73 mmol) and 12.2 mL of triethylamine (87.6 mmol) in 200 mL of THF was cooled to -78 °C. To this solution was added 9.8 mL of trimethylacetyl chloride (80 mmol) and the reaction was warmed to 0 °C and stirred for 1 hr. This solution
 10 was recooled to -78 °C. (Solution #1) Meanwhile, to a solution of 14.2 gm of (S)-(-)-4-benzyl-2-oxazolidinone in 200 mL of THF at -78 °C, was added 50 mL of a 1.6M solution of n-butyl lithium in THF (80 mmol) over 20 min. (Solution #2) After stirring for 20 min., this solution (#2) was slowly cannulated into the solution (#1) containing the mixed anhydride at -78
 15 °C. This reaction mixture was allowed to warm to room temperature and stirred for 2 hr. The reaction was quenched with a solution of saturated NH₄Cl and extracted with ethyl acetate. The combined organic fractions were washed with water, sat. NaHCO₃ solution, sat. NaCl solution, dried over MgSO₄, filtered and concentrated. The crude
 20 product was purified by chromatography (silica, hexanes : ethyl; acetate = 4 : 1) to give the title compound.

Step 2 3-(2-Phenyl-2-(S)-allylacetyl)-4-(S)-benzyl-2-oxazolidinone

To a solution of 15.5 gm of 3-phenylacetyl-4-(S)-benzyl-2-oxazolidinone (52 mmol) in 100 mL of THF at -78 °C in a nitrogen
 25 atmosphere was added dropwise 63 mL of a 1M solution of lithium

bis(trimethylsilyl)azide (63 mmol). The reaction mixture was warmed to 0 °C over the period of 1 hr and then recooled to -78 °C. To the reaction mixture was added 5.3 mL of allyl iodide (57 mmol). The reaction mixture was slowly warmed to room temperature and stirred for 12 hr.

- 5 The reaction was quenched with saturated NH₄Cl solution and the mixture was extracted with ethyl acetate. The combined organic fractions were washed with solutions of saturated NaHCO₃ and NaCl, dried over MgSO₄, filtered and concentrated. The residue was purified by chromatography (silica, 2:1 hexanes:ethyl acetate) to give the title
10 compound.

¹H NMR (CDCl₃) δ 2.5 (m, 1H), 2.8 (ABq, 1H), 3.00 (m, 1H), 3.35 (d, 1H), 4.1 (m, 2H), 5.1 (d, 1H), 5.1.5 (d, 1H), 5.2(t, 1H), 5.8 (m, 1H), 7.3 (m, 10H).

Step 3 2-(S)-allyl-2-phenethanol

- 15 To a suspension of 4.57 gm of LAH (115 mmol) in 150 mL of THF at 0 °C was added a solution of 19.2 gm of 3-(2-Phenyl-2-(S)-allylacetyl)-4-(S)-benzyl-2-oxazolidinone (57 mmol) in 100 mL of THF. The reaction mixture was stirred for 12 hr at room temperature. The reaction was quenched with 10 mL of water, and the pH was lowered to
20 pH=1 with a 2N solution of H₂SO₄. The mixture was extracted with ethyl acetate and the combined organic fractions were dried over MgSO₄, filtered and concentrated. The title compound was used without further purification. ¹H NMR (CDCl₃) δ 1.4 (br s, 1H), 2.45 (m, 2H), 2.9 (m, 1H), 3.8 (m, 2H), 5.0 (d, 1H), 5.05 (d, 1H), 5.75 (m, 1H), 7.3 (m, 5H).

25

Step 4 2-(S)-allyl-2-phenethyl bromide

- To a solution of 18.5 gm of triphenylphosphine (70.6 mmol) in 200 mL of anhydrous ether at room temperature was added 23.3 gm of carbon tetrabromide (70.6 mmol). After stirring for 30 min, a solution of
30 6.76 gm of 3-(2-Phenyl-2-(S)-allylethanol (44.1 mmol) in 50 mL of ether

was added dropwise and the reaction was allowed to stir for 12 hr. The reaction mixture was filtered and the filtercake was washed with ether. The combined ether fractions were washed with solutions of saturated NaHCO₃ and NaCl, dried over MgSO₄, filtered and concentrated. The residue was purified by chromatography (silica, 4:1, hexanes: ethyl acetate) to give the title compound.

¹H NMR (CDCl₃) δ 2.5 (m, 1H), 2.65 (m, 1H), 3.05 (m, 1H), 3.6 (m, 2H), 5.0 (d, 1H), 5.05 (d, 1H), 5.65 (m, 1H), 7.3 (m, 5H).

10 Step 5 2-(S)-allyl-2-phenethyl amine

To a solution of 8.6 gm of 2-(S)-allyl-2-phenethyl bromide (37 mmol) in 60 mL of DMF was added 13.7 gm of potassium phthalimide (74 mmol, Aldrich) and the reaction mixture was heated at 78 °C for 12 hr. The reaction was quenched with water and the mixture was extracted with ethyl acetate. The combined organic fractions were washed with solutions of saturated NaHCO₃ and NaCl, dried over MgSO₄, filtered and concentrated. The residue was purified by chromatography (silica, 9:1, hexanes: ethyl acetate) to give 4.77 gm of the phthalimide intermediate. A solution of the phthalimide and 2.2 mL of hydrazine hydrate (70.5 mmol) in 50mL of ethanol was stirred for 3 hr. The reaction mixture was diluted with water, made acidic with HCl and washed with ethyl acetate. The aqueous layer was then made basic with a solution of 1N KOH and extracted with ether. The combined ether fractions were washed with solutions of saturated NaHCO₃ and NaCl, dried over MgSO₄, filtered and concentrated to give the title compound, which was used without further purification.

¹H NMR (CDCl₃) δ 1.9 (br s, 2H), 2.4 (m, 2H), 2.7 (m, 1H), 2.8 (m, 1H), 3.0 (m, 1H), 4.95 (d, 1H), 5.0 (d, 1H), 5.7 (m, 1H), 7.3 (m, 5H).


30 Step 5 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-allyl-4-phenylbutane

The title compound was prepared from 2-(S)-allyl-2-phenethyl amine and 4 methoxybenzoyl chloride according to procedures described in Example 2.

- ^1H NMR (CDCl_3) δ 2.45 (m, 2H), 3.0 (m, 1H), 3.45 (m, 1H), 3.55 (s, 3H),
5 4.05 (m, 1H), 4.95 (d, 1H), 5.05 (d, 1H), 5.7 (m, 1H), 6.8 (d, 1H), 7.0 (t, 1H),
7.3 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 296 (M^+),
 α_D (CHCl_3) = -14.3

EXAMPLE 17

- 10 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-allyl-4-phenylbutane




The title compound was prepared according to procedures described in Example 16, with the exception that (R)-(-)-4-benzyl-2-oxazolidinone was used as the chiral auxiliary.

- 15 ^1H NMR (CDCl_3) same as for Example 16, Step 5. Mass Spectrum m/e 296 (M^+), α_D (CHCl_3) = +14.27.

EXAMPLE 18

- 20 1-(2-methoxyphenyl)-1-oxo-2-aza-(S)-4-i-butyl-4-phenylbutane



Step 1 2-(S)-i-butyl-2-phenethanol

The title compound was prepared according to procedures described in Example 16, Steps 1-3.

5 Step 2 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-i-butyl-4-phenylbutane


2-(S)-i-butyl-2-phenethanol was converted to the title compound according to procedures described in Example 11, Steps 2 and 3.

10 ^1H NMR (CDCl_3) δ 0.9 (t, 6H), 1.5 (m, 3H), 1.65 (m, 2H), 3.0 (m, 1H), 3.35 (m, 1H), 3.55 (s, 3H), 6.8 (d, 1H), 7.0 (t, 1H), 7.3 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 312 (M^+), α_D (CHCl_3) = -21.4

EXAMPLE 19

15

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-n-propyl-4-phenylbutane




The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-3-(S)-allyl-4-phenylbutane according to procedures described in Example 6.

20 ^1H NMR (CDCl_3) δ 0.9 (t, 3H), 1.3 (m, 2H), 1.7 (m, 3H), 2.9 (m, 1H), 3.4 (m, 1H), 3.6 (s, 3H), 4.05 (m, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.35 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 298 (M^+), α_D (CHCl_3) =
25 -22.4

EXAMPLE 20

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-(3-hydroxypropyl)-4-phenylbutane




- 5 The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-allyl-4-phenylbutane according to procedures described in Example 7.

^1H NMR (CDCl_3) δ 1.5 (m, 2H), 1.6 (br s, 1H), 1.85 (m, 2H) 2.9 (m, 1H),
 3.45 (m, 1H), 3.6 (s, m, 5H), 4.00 (m, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.35 (m,
 10 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 314 (M^+), α_D
 (CHCl_3) = -19.6

EXAMPLE 21

- 15 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-((3-allylcarbamoyloxy)propyl)-4-phenylbutane




To solution of 0.2 gm of 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-(3-hydroxypropyl)-4-phenylbutane (0.638 mmol) in 5 mL of

dichloromethane and 0.176 mL of triethylamine (1.27 mmol) and 0.116 gm of DMAP (0.96 mmol) was added 0.19 gm of 4-nitrophenyl chloroformate and the reaction was stirred at rt. After 3 hr, the reaction mixture was concentrated and purified by chromatography (silica, hexane : ethyl acetate = 1: 1) to give the 4-nitrophenyl carbonate intermediate. The carbonate was redissolved in 2 mL of dichloromethane and to it was added 0.18 mL of allylamine (2.4 mmol) and the reaction was stirred for 12 hr at rt. The reaction mixture was washed with saturated NaHCO₃, 3N HCl and NaCl, dried over MgSO₄, filtered and concentrated. The residue was purified by chromatography (silica, silica, hexane : ethyl acetate = 1: 1) to give the title compound.

¹H NMR (CDCl₃) δ 1.6 (m, 2H), 1.7 (m, 1H), 1.85 (m, 1H) 2.9 (m, 1H), 3.4 (m, 1H), 3.6 (s, 3H), 3.8 (m, 1H), 4.1 (m, 4H), 4.95 (br s, 1H), 5.1 (d, 1H), 5.2 (d, 1H), 5.8 (br s, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 397 (M⁺1), α_D (CHCl₃) = -21.4

EXAMPLE 22

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-((3-allyloxycarbonyloxy)propyl)-4-phenylbutane



The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-(3-hydroxypropyl)-4-phenylbutane


according to procedures described in Example 21 with the exception that allylchloroformate was used.

^1H NMR (CDCl_3) δ 1.6 (m, 1H), 1.65 (m, 1H), 1.7 (m, 1H), 1.9 (m, 1H) 2.9 (m, 1H), 3.4 (m, 1H), 3.6 (s, 3H), 4.0 (m, 1H), 4.1 (m, 2H), 4.6 (d, 2H), 5.25 (d, 1H), 5.35 (d, 1H), 5.9 (m, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 398 (M^+), α_D (CHCl_3) = -12.6

EXAMPLE 23

10

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-n-propyl-4-phenylbutane




The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-allyl-4-phenylbutane (Example 17) according to procedures described in Example 6.

^1H NMR (CDCl_3) δ 0.9 (t, 3H), 1.3 (m, 2H), 1.7 (m, 3H), 2.9 (m, 1H), 3.4 (m, 1H), 3.6 (s, 3H), 4.05 (m, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.35 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 298 (M^+), α_D (CHCl_3) = +22.7

20

EXAMPLE 24

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-(3-hydroxypropyl)-4-phenylbutane




The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-allyl-4-phenylbutane according to procedures described in Example 7.

- 5 ^1H NMR (CDCl_3) δ 1.5 (m, 2H), 1.6 (br s, 1H), 1.85 (m, 2H) 2.9 (m, 1H), 3.45 (m, 1H), 3.6 (s, m, 5H), 4.00 (m, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.35 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 314 ($M+1$), α_D (CHCl_3) = +18.2

10

EXAMPLE 25

1-(2-methoxyphenyl)-1-oxo-2-aza-3-(R)-((3-allylcarbamoyloxy)propyl)-4-phenylbutane



15

The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-(3-hydroxypropyl)-4-phenylbutane according to procedures described to Example 21.


^1H NMR (CDCl_3) δ 1.6 (m, 2H), 1.7 (m, 1H), 1.85 (m, 1H) 2.9 (m, 1H), 3.4 (m, 1H), 3.6 (s, 3H), 3.8 (m, 1H), 4.1 (m, 4H), 4.95 (br s, 1H), 5.1 (d, 1H), 5.2

(d, 1H), 5.8 (br s, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 397 (M⁺1), α_D (CHCl₃) = +19.8

EXAMPLE 26

5

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-((3-allyloxycarbonyloxy)propyl)-4-phenylbutane




The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4-(R)-(3-hydroxypropyl)-4-phenylbutane according to procedures described in Example 21 with the exception that allylchloroformate was used.

¹H NMR (CDCl₃) δ 1.6 (m, 1H), 1.65 (m, 1H), 1.7 (m, 1H), 1.9 (m, 1H) 2.9 (m, 1H), 3.4 (m, 1H), 3.6 (s, 3H), 4.0 (m, 1H), 4.1 (m, 2H), 4.6 (d, 2H), 5.25 (d, 1H), 5.35 (d, 1H), 5.9 (m, 1H), 6.85 (d, 1H), 7.05 (t, 1H), 7.3 (m, 6H), 7.8 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 398 (M⁺1), α_D (CHCl₃) = +12.6.

20

EXAMPLE 27

1-(2-methoxyphenyl)-1-oxo-2-aza-4-hydroxyethyl-4-(4-oxy-3-oxo-pentanyl)-4-phenylbutane



Step 1 4-cyano-4-phenylcyclohexanone ethyleneglycol ketal

A solution of 20 gm of 4-cyano-4-phenylcyclohexanone (Aldrich, 100.5 mmol), 60 ml of ethylene glycol, and 1.10 gm of TsOH in 65 ml of toluene was refluxed for 12 hr. The water formed during the reaction was removed via a Dean-Stark distillation receiver. The reaction mixture was concentrated under reduced pressure to remove the solvent and the residue was poured into 200 ml of ether. It was washed with water (20 ml x 3), dried over MgSO_4 and concentrated. The residue was purified by recrystallization with hexanes: ether to afford the title compound as a white solid.

Step 2 4-aminomethyl-4-phenylcyclohexanone ethyleneglycol ketal

To a suspension of 10.00 g (41.1 mmol) of 4-cyano-4-phenylcyclohexanone ethylene glycol ketal in 50 ml of dry THF was slowly added 61.65 ml of lithium aluminum hydride (1.0 M in THF, 61.65 mmol) and the reaction mixture was refluxed for 3 hr. When TLC showed no starting material the reaction mixture was cooled to 0 °C and quenched with 4 ml of 4N NaOH at 0 °C. The reaction mixture was filtered through a plug of Na_2SO_4 and concentrated to give the title compound as a colorless oil.


Step 3 4-Phenyl-4-(3-(2-methoxyphenyl)-3-oxo-2-azaprop-1-yl)cyclohexanone

To a solution of 6.06 g (24.5 mmol) of 4-aminomethyl-4-phenylcyclohexanone ethyleneglycol ketal and 6.50 ml of triethylamine

(49.0 mmol) in 60 ml of methylene chloride was added 5.44 g (31.9 mmol) of o-anisoyl chloride at 0 °C. The reaction mixture was stirred for 3 hr and was poured into 200 ml of ether. It was washed with aq NaHCO₃, dried over MgSO₄ and concentrated. Then 150 ml of THF and 50 ml of 2N HCl was added into the residue. The reaction mixture were stirred for 3 hr at 40 °C. Then it was poured into 200 ml of ether. The organic layer was washed with aq NaHCO₃, dried over MgSO₄ and concentrated. The residue was purified by chromatography (silica, hexanes: ethyl acetate, 1:1) to afford the compound as a solid.

10

Step 4 5-phenyl-4-(3-(2-methoxyphenyl)3-oxo-2-aza-propyl)-1-oxo-2-oxocycloheptane



15 A solution of 0.044 gm of 4-phenyl-4-(3-(2-methoxyphenyl)3-oxo-2-aza-propyl)-cyclohexanone (Lancaster Chemical Co., 0.13 mmol) and 0.15 gm of 3-chloro peroxybenzoic acid (0.87 mmol) in 10 mL of CH₂Cl₂ was stirred at rt for 72 hr and was filtered through a plug of silica gel. The residue was purified by HPLC (Waters RCM 25x10, with
20 CH₃CN:tBuOMe:Hexane (1:4:5)/Hexane = 6.75/6.0 mL/min) to give the title compound as a white solid.

¹H NMR (CDCl₃) δ 8.16 (d of d, 1H, J = 1.3, 7.7 Hz); 7.63 (t, 1H, J = 5.5 Hz), 7.47 (t, 2H, J = 7.6 Hz), 7.41 (t of d, 1H, J = 1.1, 8.2 Hz), 7.34-7.39 (m, 3H), 4.36 (d of d, 1H, J = 7.1, 13.3 Hz), 4.13 (d of d, 1H, J = 9.1, 13.2 Hz), 3.64 -
25 3.73 (m, 2H), 3.62 (s, 3H), 2.72 (d of d, 1H, J = 9.1, 14.2 Hz), 2.54 - 2.61 (m,


2H), 2.45 (d of d, 1H, J = 14.2, 8.9 Hz), 2.11 (d of d, 1H, J = 9.1, 16 Hz), 2.02 (d of d, 1H, J = 11.6, 14.4 Hz); Mass Spectrum (CI) m/e 354 (M+1)

Step 5 1-(2-methoxyphenyl)-1-oxo-2-aza-4-hydroxyethyl-4-(4-oxy-3-oxo-pentanyl)-4-phenylbutane

5 A solution of 0.0081 gm of 5-phenyl-4-(3-(2-methoxyphenyl)3-oxo-2-aza-propyl)-1-oxy-2-oxocycloheptane (0.023 mmol) and 0.0065 gm of potassium carbonate (0.047 mmol) in 4 mL of methanol was stirred at rt for 14 hr and solvent was removed. The
10 residue was dissolved in EtOAc and filtered through a plug of silica gel to give the title compound.
¹H NMR (CDCl₃) δ 8.19 (d of d, 1H, J = 1.7, 7.8 Hz); 7.99 (bs 1H), 7.37 - 7.45 (m, 5H), 7.27 - 7.29 (m, 1H), 7.07 (t, 1H, J = 7.7 Hz), 6.90 (d, 1H, J = 8.5 Hz), 4.05 (d of d, 1H, J = 7.1, 13.8 Hz), 3.80 (d of d, 1H, J = 5.5, 14.0 Hz), 3.60 -
15 3.73 (m, 2H), 3.66 (s, 3H), 3.62 (s, 3H), 1.98 - 2.25 (m, 7H); Mass Spectrum (CI) m/e 386 (M+1)

EXAMPLE 28

20 1-(2-methoxyphenyl)-1-oxo-2-aza-4-hydroxyethyl-4-(8-phenyl-4-oxy-3-oxo-octanyl)-4-phenylbutane



A solution of 0.01 gm of 5-phenyl-4-(3-(2-methoxyphenyl)3-oxo-2-aza-propyl)-1-oxy-2-oxocycloheptane (0.028 mmol), 0.005 mg
25 potassium carbonate (0.036 mmol) and (0.042 gm, 0.28 mmol) of 4-phenyl-1-butanol in 1.0 mL of DMF was heated to 80 °C for 14 hr and

solvent was removed. The residue was dissolved in EtOAc and filtered through a plug of silica gel. It was purified by HPLC (Waters RCM 25x10, with CH₃CN:tBuOMe:Hexane (1:4:5)/Hexane = 6.75/6.0 mL/min) to give the title compound.


¹H NMR (CDCl₃) δ 8.19 (d, 1H, J = 7.8 Hz), 8.01 (t, 1H, J = 5.8 Hz), 7.16 - 7.44 (m, 11H), 7.06 (t, 1H, J = 7.5 Hz), 6.89 (d, 1H, J = 8.5 Hz), 4.01 - 4.04 (m, 3H), 3.80 (d of d, 1H, J = 5.5, 13.8 Hz), 3.62 - 3.75 (m, 2H), 3.65 (s, 3H), 2.79 (bs, 1H), 2.63 (t, 2H, J = 7.0 Hz), 1.98 - 2.25 (m, 6H), 1.63 - 1.67 (m, 4H); Mass Spectrum (CI) m/e 504 (M+1).

10

EXAMPLE 29

1-(2-methoxyphenyl)-1-oxo-2-aza-4-hydroxyethyl-4-(4-oxy-3-oxo-hept-6-enyl)-4-phenylbutane

15




The title compound was prepared from 30.6 mg 5-phenyl-4-(3-(2-methoxyphenyl)3-oxo-2-aza-propyl)-1-oxy-2-oxocycloheptane as in Example 28 with the exception that allyl alcohol was used and the reaction was stirred at 40 °C for 14 hr.

¹H NMR (CDCl₃) δ 8.16 (d of d, 1H, J = 1.4, 7.8 Hz); 8.01 (t, 1H, J = 5.5 Hz), 7.37 - 7.42 (m, 5H), 7.25 - 7.28 (m, 1H), 7.03 (t, 1H, J = 7.5 Hz), 6.87 (d, 1H, J = 8.2 Hz), 5.85 (m, 1H), 5.28 (d of d, 1H, J = 1.3, 17.2Hz), 5.19 (d, 1H, J = 11.4 Hz), 4.50 (d, 2H, J = 5.7 Hz), 4.01 (d of d, 1H, J = 6.8, 13.7 Hz), 3.77 (d of d, 1H, J = 5.5, 13.7 Hz), 3.59 - 3.72 (m, 2H), 3.64 (s, 3H), 2.96 (bs, 1H), 1.97 - 2.27 (m, 6H); Mass Spectrum (CI) m/e 412 (M+1)

EXAMPLE 30


1-(2-methoxyphenyl)-1-oxo-2-aza-4-acetoxyethyl-4-(4-oxy-3-oxo-hept-6-enyl)-4-phenylbutane



A solution of 0.0095 gm of 1-(2-methoxyphenyl)-1-oxo-2-aza-4-hydroxyethyl-4-(4-oxy-3-oxo-hept-6-enyl)-4-phenylbutane (0.023 mmol), 0.0018 gm of DMAP, 0.05 mL of Ac₂O and 0.1 mL of pyridine in 2.0 mL of THF was stirred at rt for 14 hr and volatiles were removed under vacuum. The residue was dissolved in CH₂Cl₂ and filtered through a plug of silica gel, purified by HPLC (Waters RCM 25x10, with CH₃CN:tBuOMe:Hexane (1:4:5)/Hexane = 6.75/6.0 mL/min) conditions) to afford the title compound. ¹H NMR (CDCl₃) δ 8.18 (d of d, 1H, J = 1.8, 7.8 Hz), 7.80 (t, 1H, J = 5.3 Hz), 7.39 - 7.42 (m, 5H), 7.27 - 7.32 (m, 1H), 7.05 (t, 1H, J = 7.8 Hz), 6.88 (d, 1H, J = 8.2 Hz), 5.83 - 5.90 (m, 1H), 5.27 (d, 1H, J = 15.2 Hz), 5.20 (d, 1H, J = 10.3 Hz), 4.52 (d, 2H, J = 5.7 Hz), 4.07 (t, 2H, J = 7.1 Hz), 3.91 (d of d, 1H, J = 6.0, 13.8 Hz), 3.83 (d of d, 1H, J = 5.5, 13.7 Hz), 3.63 (s, 3H), 2.29 - 2.38 (m, 2H), 2.11 - 2.18 (m, 4H), 1.96 (s, 3H); Mass Spectrum (CI) m/e 454 (M+1).

EXAMPLE 31


1-(2-methoxyphenyl)-1-oxo-2-aza-4-hydroxyethyl-4-(4-aza-3-oxo-pentanyl)-4-phenylbutane



The title compound was prepared (from 27.1 mg of 5-phenyl-4-(3-(2-methoxyphenyl)3-oxo-2-aza-propyl)-1-oxy-2-oxocycloheptane (0.077 mmol) according to procedures described in Example 28 with the exception that the reaction was carried out in a solution of 2M methylamine in THF. ¹H NMR (CDCl₃) δ 8.25 (t, 1H, J = 5.7 Hz), 8.12 (d, 1H, J = 6.8 Hz); 7.42 (t, 1H, J = 7.0 Hz), 7.32 - 7.36 (m, 4H), 7.21 - 7.23 (m, 1H, J = 6.8 Hz); 7.04 (t, 1H, J = 7.3 Hz), 6.89 (d, 1H, J = 8.2 Hz), 6.60 (q, 1H, J = 4.1 Hz), 3.89 (d, 2H, J = 6.2 Hz), 3.68 (s, 3H), 3.58 - 3.62 (m, 1H), 3.41 - 3.46 (m, 1H), 3.25 (bs, 1H), 2.66 (d, 4.8 Hz), 1.89 - 2.18 (m, 6H); Mass Spectrum (CI) m/e 385 (M+1)

EXAMPLE 32

15 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diphenylbutane



To a solution of 0.106 gm (0.58 mmol) of 1-amino diphenylmethane in 5 mL of dichloromethane and 0.1 mL of pyridine (1.16 mmol) at 0 °C was added 0.198 gm of 2-methoxybenzoyl chloride (0.172 mL, 1.16 mmol). The reaction was allowed to warm to rt and was stirred for 4 hr. The reaction mixture was concentrated, 5mL of heptane were added and the mixture was evaporated to dryness this being repeated one more time. The residue was purified by HPLC (RCM

SepPak, silica 25x100, 1.25% CH₃CN, 5% tBuOCH₃, 93.75% hexane) to give the title compound.

¹H NMR (500MHz, CDCl₃) δ 3.95 (s, 3H), 6.58 (d, 1H), 7.01 (d, 1H), 7.06 (t, 1H), 7.25 - 7.50 (m, 9H), 8.2 (d, 1H), 8.7 (d, 1H); ¹³C NMR (CDCl₃) δ 55.82,


5 57.15, 164.16; Mass Spectrum m/e 318 (M⁺1).

The following examples 33 to 37 were prepared as described in Examples 3 and 2.

10

EXAMPLE 33

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(2-methoxyphenyl)-butane




15 ¹H NMR (CDCl₃) δ 0.8 (t, 6H), 1.8 (m, 2H), 2.0 (m, 2H), 3.5 (s, 3H), 3.85 (s, 3H), 4.0 (d, 2H), 6.83 (d, 1H), 6.94 (d, 1H), 6.98 (t, 1H), 7.04 (t, 1H), 7.28 (m, 1H), 7.31 (m, 1H), 7.37 (m, 1H), 7.6 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 342 (M⁺1).

20

EXAMPLE 34

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(3-methoxyphenyl)-butane




¹H NMR (CDCl₃) δ 0.8 (t, 6H), 1.78 (m, 4H), 3.6 (s, 3H), 3.76 (d, 2H), 3.83 (t, 3H), 6.8 (dd, 1H), 6.86 (d, 1H), 6.97 (t, 1H), 7.02 (m, 1H), 7.04 (m, 1H), 7.32 (t, 1H), 7.4 (m, 1H), 7.66 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e

5 342 (M⁺1).

EXAMPLE 35

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(4-methoxyphenyl)-butane




10

¹H NMR (CDCl₃) δ 0.8 (t, 6H), 1.76 (m, 4H), 3.6 (s, 3H), 3.77 (d, 2H), 3.83 (t, 3H), 6.88 (d, 1H), 6.94 (m, 2H), 7.05 (t, 1H), 7.32 (m, 2H), 7.4 (m, 1H), 7.65 (br s, 1H), 8.2 (dd, 1H). Mass Spectrum m/e 342 (M⁺1).

15

EXAMPLE 36


1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(2,4-dimethylphenyl)-butane



¹H NMR (CDCl₃) δ 0.8 (t, 6H), 2.2 (m, 4H), 3.4 (s, 3H), 4.2 (d, 2H), 6.78 (d, 1H), 7.02 (t, 1H), 7.36 (dt, 1H), 7.5 (m, 3H), 7.56 (d, 1H), 7.7 (br s, 1H), 7.8 (d, 1H), 7.92 (dd, 1H), 8.2 (dd, 1H), 8.55 (s, 1H). Mass Spectrum m/e 340 (M+1).

EXAMPLE 37

10 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(naphth-1-yl)-butane




¹H NMR (CDCl₃) δ 0.8 (t, 6H), 1.76 (m, 4H), 2.3 (s, 3H), 2.5 (s, 3H), 3.5 (s, 3H), 3.9 (d, 2H), 6.84 (d, 1H), 7.02 (m, 1H), 7.05 (t, 1H), 7.25 (d, 1H), 7.39 (m, 1H), 7.7 (br s, 1H), 8.22 (dd, 1H). Mass Spectrum m/e 362 (M+1).


15

EXAMPLE 38

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(4-acetylphenyl)-butane



Step 1 **2-(4-Tributyltinphenyl)acetonitrile**




5

To a solution of 10 gm of 4-bromophenylacetonitrile (50 mmol) in 400 mL THF at -90°C was added 100 mL of t-butyl lithium (1 M in hexanes 170 mmol). To this was added a solution of 19.5 gm of tributyltin chloride (60 mmol) in 10 mL THF and the reaction mixture was allowed to warm to -70 °C. The reaction mixture was stirred for 30 min and then poured into 400 mL water. The reaction mixture was extracted with 500 mL of 1:1 ether-hexanes and the layers were separated. The organic layer was washed with water and brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by chromatography (silica, 30% ether-hexanes) to afford the title compound as a colorless oil.

¹H NMR (CDCl₃) δ 0.80 (t, 9H), 1.06 (m, 6H), 1.32 (m, 6H), 1.52 (m, 6H), 3.73 (s, 2H), 7.28 (d, 2H), 7.47 (d, 2H).

20


Step 2 **2-(4-Tributyltinphenyl)-2,2-diethylacetonitrile**



To a solution of 10.1 gm of 2-(4-tributyltinphenyl)acetonitrile (Step 1, 25 mmol) in 250 mL of tetrahydrofuran at -20 °C was added 75 mL
5 t-BuOK (1M in t-butanol, 75 mmol), and the reaction mixture was stirred for 5 min. To the reaction mixture was then added 10.1 gm of ethyl iodide (65 mmol) and stirring was continued for 1 hr as the temperature was warmed to rt. The reaction mixture was then partitioned between
10 NH₄Cl (sat) and ether. The organic fraction was washed with H₂O, brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by chromatography (silica, 5-15% ether-hexane) to afford the title compound as a colorless oil.
¹H NMR (CDCl₃) δ 0.80 - 0.95 (m, 15H), 1.06 (m, 6H), 1.32 (m, 6H), 1.52 (m, 6H), 1.82- 2.05 (m, 4H), 7.28 (m, 2H), 7.47 (m, 2H).

15


Step 3 2-(4-tributyltinphenyl)-(2,2-diethyl)ethylamine



To a solution of 0.3 gm of 2-(4-Tributyltinphenyl)-2,2-
20 diethylacetonitrile (Step 2, (1.1 mmol)) in 5 mL of MeOH and 0.5 mL of THF was added 0.1 gm (0.78 mmol) of CoCl₂. After stirring for for 5 min., 0.065 of NaBH₄ (1.75 mmol) was added portionwise over 3 hrs. When the reaction was complete (TLC 90:10:1 CHCl₃-MeOH-NH₄OH), it was partitioned between NH₄Cl (sat) and ether. The organic fraction


was washed with water and brine, dried over Na_2SO_4 , filtered and concentrated. The residue was used without further purification.

Step 4
5 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(4-tributyltinphenyl)-butane



10 To a solution of 2-(4-tributyltinphenyl)-(2,2-diethyl)ethylamine (~1 mmol) in 5 mL dichloromethane was added 0.12 mL of pyridine (3 mmol), 0.024 gm of DMAP (0.2 mmol) and 0.34 gm of o-anisoyl chloride (2 mmol). After stirring for 1 hr, the reaction mixture was partitioned between ethyl ether and water. The organic fraction
15 was washed with water and brine, dried over NaSO_4 , filtered and concentrated. The residue was purified by chromatography (silica, 20-25% ether-hexane) to afford the title compound as a colorless oil.
 ^1H NMR (CDCl_3) δ 0.82 (t, 6H), 0.91 (t, 9H), 1.07 (t, 6H), 1.36(m, 6H), 1.56(m, 6H), 1.79(q, 4H), 3.62 (s, 3H), 3.77 (d, 2H), 6.84 (d, 1H), 7.04 (t, 1H),
20 7.35 -7.40 (m, 3H), 7.49 (d, 2H) 7.68 (br s, 1H), 8.21 (dd, 1H).


Step 5 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(4-acetylphenyl)-butane



To a solution of 0.24 gm of 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(4-tributyltinphenyl)butane (0.04 mmol) in 1 mL of
 5 dichloromethane was added 4 mg triethylamine (0.04 mmol), 0.01 gm of K_2CO_3 , 0.002 gm, $Pd_2(dba)_3 \cdot CHCl_3$ (5 mole%) and 0.004 gm of acetyl chloride (0.05 mmol), the reaction mixture was stirred for 4 hr at ambient temperature. The reaction mixture was partitioned between water and ether and the organic fraction was dried over $MgSO_4$, filtered
 10 and concentrated. The residue was purified by HPLC (Waters RCM, Nova-Pak silica, 8 mm X 10 cm, using a mixture of 2:1 (6:3:1 hexane-methyl *tert*-butyl ether-acetonitrile:hexane) to afford the title compound. 1H NMR ($CDCl_3$) δ 0.82 (t, 6H), 1.83 (m, 4H), 2.63 (s, 3H), 3.60 (s, 3H), 3.84 (d, 2H), 6.86 (d, 1H), 7.06 (t, 1H), 7.40(m, 1H), 7.52 (d, 2H) 7.65 (m, 1H),
 15 7.99 (m, 2H), 8.22 (dd, 1H) Mass Spectrum (CI) m/e 354 (M+1)

EXAMPLE 39

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(4-bromophenyl)-butane



20


To a solution of 0.24 of 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(4-tributyltinphenyl)-butane (0.04 mmol) in 1 mL dichloromethane was added 0.008 gm triethylamine (0.08 mmol) and 0.08 mL of a 1M solution of bromine in dichloromethane. After stirring

for 15 min., the reaction was quenched with water and a drop of sat. NaHSO₃. The organic fraction was dried over MgSO₄, filtered and concentrated. The residue was purified by HPLC (Waters RCM, Nova-Pak silica, 8 mm X 10 cm) using a mixture of 2:1 (6:3:1 hexane-methyl
5 *tert*-butyl ether-acetonitrile:hexane) to afford the title compound.
¹H NMR (CDCl₃) δ 0.81 (t, 6H), 1.76 (m, 4H), 3.62 (s, 3H), 3.77 (d, 2H), 6.87 (d, 1H), 7.06 (t, 1H), 7.29 (d, 2H), 7.40(m, 1H), 7.52 (d, 2H) 7.63 (br s, 1H), 8.22 (dd, 1H). d Mass Spectrum (CI) m/e 390,392 (M+1).

10

EXAMPLE 40

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(3-acetylphenyl) butane




15

The title compound was prepared from 3-bromophenyl acetonitrile according to procedures described in Example 38.

¹H NMR (CDCl₃) δ 0.81 (t, 6H), 1.85 (m, 4H), 2.63 (s, 3H), 3.59 (s, 3H), 3.84 (d, 2H), 6.86 (d, 1H), 7.06 (t, 1H), 7.40(m, 1H), 7.50 (d, 1H) 7.64 (m, 2H), 7.86 (m, 1H), 8.03 (m, 1H), 8.22 (dd, 1H) Mass Spectrum (CI) m/e 354
20 (M+1)

EXAMPLE 41

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(3-bromophenyl)-butane




The title compound was prepared from 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(3-tributyltinphenyl)-butane according to procedures described in Example 39.

¹H NMR (CDCl₃) δ 0.82 (t, 6H), 1.78 (m, 4H), 3.65 (s, 3H), 3.77 (d, 2H), 6.88 (d, 1H), 7.06 (t, 1H), 7.28 (m, 1H), 7.35 - 7.45 (m, 3H), 7.55 (m, 1H) 7.63 (br s, 1H), 8.22 (dd, 1H). d Mass Spectrum (CI) m/e 390,392 (M+1)

10

EXAMPLE 42

1-(2-methoxy-5-bromophenyl)-1-oxo-2-aza-4,4-diethyl-4-(3-bromophenyl)butane



15

To a solution of 0.024 gm of 1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-(3-tributyltinphenyl)-butane (0.04 mmol) in 1 mL of dichloromethane was added 0.008 gm of triethylamine (0.08 mmol) and 1.2 mL of a 1M solution of bromine in dichloromethane. After stirring for 45 min. the reaction mixture was partitioned into water and a drop of sat. NaHSO₃. The organic fraction was dried over MgSO₄, filtered and concentrated. The residue was purified by HPLC (Waters RCM, Nova-


20

Pak silica, 8 mm X 10 cm) using a mixture of 2:1 (6:3:1 hexane-methyl *tert*-butyl ether-acetonitrile:hexane) to afford the title compound.

^1H NMR (CDCl_3) δ 0.82 (t, 6H), 1.77 (m, 4H), 3.63 (s, 3H), 3.75 (d, 2H), 6.77 (d, 1H), 7.28 (t, 1H), 7.35 (d, 1H), 7.42 (d, 1H), 7.49 (dd, 1H), 7.50-7.56 (br m, 2H), 8.32 (dd, 1H). d Mass Spectrum (CI) m/e


EXAMPLE 43

10 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(phenyl)-butane



Step 1 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(4-tributyltinphenyl)-butane

15



To a solution of 2-(4-tributyltinphenyl)-(2,2-diethyl) ethylamine (Example 38, Step 3, ~1 mmol) in 5 mL dichloromethane was added 0.12 mL pyridine (3 mmol) , 0.024 gm of DMAP (0.2 mmol) and 0.36 gm of 7-dihydrobenzofuranyl chloride (2 mmol) and the reaction mixture stirred for 1 hr. The mixture was partitioned between ethyl ether and water and the organic fraction was washed with water and brine, dried over NaSO_4 filtered and concentrated. The residue was purified by chromatography (silica, 20-25% ether-hexane) to afford the

title compound as a colorless oil. ^1H NMR (CDCl_3) δ 0.82 (t, 6H), 0.91 (t, 9H), 1.07 (t, 6H), 1.36(m, 6H), 1.56(m, 6H), 1.79(q, 4H), 3.17(t, 2H), 3.77 (d, 2H), 4.43 (t, 2H), 6.92 (t, 1H), 7.32 -7.40 (m, 3H), 7.50 (d, 2H) 7.69 (br s, 1H), 7.91 (dd, 1H)

5


Step 2 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(phenyl)-butane

A solution of 0.025 gm 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(4-tributyltinphenyl)-butane (0.04 mmol) in 1 mL dichloromethane and 0.1 mL of 1M HCl in ether was stirred for 15 min. The reaction mixture was partitioned between 5 mL ether and 2 mL of sat. Na_2CO_3 and organic fraction was washed with water and brine, dried over NaSO_4 filtered and concentrated. The residue was purified by HPLC (Waters RCM, Nova-Pak silica, 8 mm X 10 cm) using a mixture of 2:1 (6:3:1 hexane-methyl *tert*-butyl ether-acetonitrile:hexane) to afford the title compound.

^1H NMR (CDCl_3) δ 0.81 (t, 6H), 1.80 (m, 4H), 3.17 (t, 2h), 3.76 (d, 2H), 4.44 (t, 2H), 6.93 (t, 1H), 7.26 (m, 2H), 7.32 - 7.42 (m, 5H), 7.9 (d, 1H) . d Mass Spectrum (CI) m/e 324 (M+1)

EXAMPLE 44

1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(phenyl)-butane




A solution of 0.12 gm of 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(phenyl)-butane (0.02 mmol) in n-propanol and 5% Pd/C was stirred at reflux for 3 hr. The reaction mixture was cooled, filtered and concentrated. The residue was purified by HPLC (Waters RCM, Nova-Pak silica, 8 mm X 10 cm) using a mixture of 2:1 (6:3:1 hexane-methyl *tert*-butyl ether-acetonitrile:hexane) to afford the title compound.

¹H NMR (CDCl₃) δ 0.85 (t, 6H), 1.86 (m, 4H), 3.17 (t, 2h), 3.86 (d, 2H), 6.79 (d, 1H), 7.26 (m, 2H), 7.32 - 7.42 (m, 5H), 7.9 (d, 1H) d Mass Spectrum (CI) m/e 326 (M+1)

EXAMPLE 45

1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(3-acetylphenyl)-butane




The title compound was prepared from 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(3-tributyltinphenyl)-butane according to procedures described in Examples 38 and 40.

¹H NMR (CDCl₃) δ 0.81 (t, 6H), 1.85 (m, 4H), 2.63 (s, 3H), 3.19 (t, 2H), 3.80 (d, 2H), 4.42 (t, 2H), 6.93 (d, 1H), 7.24- 7.36 (m, 3H), 7.48 (t, 1H) 7.63 (m, 1H), 7.86 (dd, 1H) 8.02 (m, 1H), Mass Spectrum (CI) m/e 366 (M+1)

EXAMPLE 46

1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(3-bromophenyl)-butane



5


The title compound was prepared from 1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(3-tributyltinphenyl)-butane according to procedures described in Example 41.

^1H NMR (CDCl_3) δ 0.81 (t, 6H), 1.79 (m, 4H), 3.20 (t, 2h), 3.73(d, 2H), 4.56 (t, 2H), 6.93 (t, 1H), 7.24 - 7.42 (m, 5H), 7.53 (m, 1H), 7.9 (d, 1H) . Mass Spectrum (CI) m/e 401,403 (m+1)

10

WHAT IS CLAIMED IS:

1. A compound of structural Formula I:



5

I

or a pharmaceutically acceptable salt, crystal form, or hydrate, wherein:

n is: 0, 1, 2 or 3;

r is: 0 or 1;

s is: 0 or 1;

10

R¹, R², R⁵, R⁶ and R⁷ are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (2) hydroxy,
- (3) (C₁-C₆)-alkyl,
- 15 (4) HO(C₁-C₆)-alkyloxy,
- (5) (C₁-C₄)-perfluoroalkyl,
- (6) (C₂-C₆)-alkenyl,
- (7) (C₂-C₆)-alkynyl,
- (8) O[(C=O)O_r]_s(C₁-C₆)-alkyl, wherein the alkyl may be cyclic
- 20 or straight-chained,
- (9) (C₁-C₆)-alkyl-S(O)_n-,
- (10) phenyl,
- (11) phenoxy,
- (12) cyano,

- (13) nitro,
(14) CO₂H,
(15) CO(C₁-C₆)-alkyl,
(16) CO₂(C₁-C₆)-alkyl,
5 (17) CONR⁸R⁹,
(18) NR⁸R⁹,
(20) (C₂-C₆)-alkenyloxy,
(21) benzyloxy,
(22) hydrogen,
10 (23) OCF₃, or
(24) R¹ and R² or R⁶ and R⁷ can be taken together when on adjacent carbons to form a fused benzo, dihydrofuranyl, furanyl, pyrrolidyl, dihydropyrrolidyl or 1,3-dioxolan group;

R³ and R⁴ are independently:

- 15 (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(2) hydroxy,
(3) HO(C₁-C₆)-alkyloxy,
(4) (C₁-C₄)-perfluoroalkyl,
(5) O(CO)CCl₃,
20 (6) (C₁-C₆)-alkyl-S(O)_n-,
(7) phenyl-(CH₂)_r-S(O)_n-,
(8) cyano,
(9) nitro,
(10) CO₂H,
25 (11) CO(C₁-C₆)-alkyl,
(12) CO₂(C₁-C₆)-alkyl,
(13) CONR⁸R⁹,
(14) NR⁸R⁹,
(15) O(CO)NR⁸R⁹,

- (16) azido,
- (17) $\text{NR}^8(\text{CO})\text{NR}^8\text{R}^9$,
- (18) hydrogen,
- (19) $(\text{C}_1\text{-C}_{10})\text{-alkyl}$, wherein alkyl includes cyclic as well as
5 acyclic groups and is unsubstituted or substituted with one
or two of the substituents selected from the group consisting
of:
- (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (b) hydroxy,
- 10 (c) oxo,
- (d) $\text{O}[(\text{C}=\text{O})\text{O}_\text{r}]_\text{s}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (e) $(\text{C}_1\text{-C}_6)\text{-alkyl-S(O)}_\text{n-}$,
- (f) aryl- $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$,
- (g) cyano,
- 15 (h) nitro,
- (i) vinyl,
- (j) NR^8R^9 ,
- (k) $\text{O}(\text{CO})\text{NR}^8\text{R}^9$,
- (l) CHO ,
- 20 (m) CO_2H ,
- (n) $\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (o) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl may be substituted
with phenyl,
- (p) CONR^8R^9 ,
- 25 (q) aryl, wherein aryl is defined as phenyl or naphthyl,
unsubstituted or substituted with one, two or three of
the substituents selected from the group consisting of:
- (a') halo, as defined above,
- (b') hydroxy,

- (c') (C₁-C₆)-alkyl,
(d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
5 (g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
(i') phenyl,
(j') phenoxy,
(k') cyano,
10 (l') nitro,
(m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
15 (q') NR⁸R⁹, and
(r) heteroaryl, wherein heteroaryl is defined as an
unsubstituted, monosubstituted, or disubstituted five
or six membered aromatic heterocycle containing
from 1 to 3 heteroatoms selected from the group
20 consisting of O, N and S and wherein the substituents
are members selected from the group consisting of:
(a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
25 (d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,

- (i') phenyl,
(j') phenoxy,
(k') cyano,
(l') nitro,
5 (m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
(q') NR⁸R⁹, and
10 (r') fused benzo or pyridyl group,
(s) heterocyclyl, wherein heterocyclyl is defined as a 3 to
7 atom cyclic, non-aromatic substituent containing
from 1 to 3 heteroatoms selected from the group
consisting of O, N, and S, said heterocycle being
15 unsubstituted or substituted with one, two or three
substituents selected from the group consisting of:
(a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
20 (d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
25 (i') phenyl,
(j') phenoxy,
(k') cyano,
(l') nitro,
(m') CO₂H,
30 (n') CO(C₁-C₆)-alkyl,

- 5 (o') CO₂(C₁-C₆)-alkyl,
 (p') CONR⁸R⁹,
 (q') NR⁸R⁹,
 (r') NR⁸CO(C₁-C₆)-alkyl,
 (s') oxo,
 (t') fused benzo, and
 (u') fused pyridyl group;
 (t) benzyl-S(O)_n-,
 (u) O[(C=O)O_R]_S(C₂-C₆)-alkenyl,
 10 (v) O[(C=O)O_R]_Saryl,
 (w) O[(C=O)O_R]_Sheteroaryl,
 (x) O(CH₂)_nheteroaryl, or
 (y) O(CH₂)_naryl;
 (20) (C₂-C₁₀)-alkenyl, wherein alkenyl is unsubstituted or
 15 substituted with one or two of the substituents selected from
 the group consisting of:
 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (b) hydroxy,
 (c) oxo,
 20 (e) (C₁-C₆)-alkyl-S(O)_n-,
 (f) phenyl-(C₁-C₆)-alkyloxy,
 (g) cyano,
 (h) nitro,
 (i) NR⁸R⁹,
 25 (j) CHO,
 (k) CO₂H,
 (l) CO(C₁-C₆)-alkyl,
 (m) CO₂(C₁-C₆)-alkyl,
 (n) CONR⁸R⁹,

- (o) aryl, wherein aryl is as defined above,
 - (p) heteroaryl, wherein heteroaryl is as defined above,
 - (q) heterocyclyl, wherein heterocyclyl is as defined above,
 - (r) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
 - 5 (s) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
 - (t) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
 - (u) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above,
 - (v) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above, and
 - (w) $O(CH_2)_n$ aryl, aryl as defined above;
- 10
- (21) (C_2-C_{10}) -alkynyl, wherein alkynyl is unsubstituted or substituted with one or two of the substituents selected from the group consisting of:
- (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 - 15 (b) hydroxy,
 - (c) oxo,
 - (d) (C_1-C_6) -alkyloxy,
 - (e) (C_1-C_6) - $S(O)_n$ -,
 - (f) phenyl- (C_1-C_6) -alkyloxy,
 - 20 (g) cyano,
 - (h) nitro,
 - (i) vinyl,
 - (j) NR^8R^9 ,
 - (k) $NR^8CO(C_1-C_6)$ -alkyl,
 - 25 (l) CHO,
 - (m) CO_2H ,
 - (n) $CO(C_1-C_6)$ -alkyl,
 - (o) $CO_2C(C_1-C_6)$ -alkyl,
 - (p) $CONR^8R^9$,

- (q) aryl, wherein aryl is as defined above,
- (r) heteroaryl, wherein heteroaryl is as defined above,
- (s) heterocyclyl, wherein heterocyclyl is as defined above,
- (t) O[(C=O)O_R]_S(C₁-C₆)-alkyl, alkyl as defined above,
- 5 (u) O[(C=O)O_R]_S(C₂-C₆)-alkenyl, as defined above,
- (v) O[(C=O)O_R]_Saryl, aryl as defined above,
- (w) O[(C=O)O_R]_Sheteroaryl, heteroaryl as defined above
- (x) O(CH₂)_nheteroaryl, heteroaryl as defined above, and
- (y) O(CH₂)_naryl, aryl as defined above,
- 10 (22) O[(C=O)O_R]_S(C₁-C₆)-alkyl, alkyl as defined above,
- (23) O[(C=O)O_R]_S(C₂-C₆)-alkenyl, as defined above,
- (24) O[(C=O)O_R]_Saryl, aryl as defined above,
- (25) O[(C=O)O_R]_Sheteroaryl, heteroaryl as defined above
- (26) O(CH₂)_nheteroaryl, heteroaryl as defined above,
- 15 (27) aryl, wherein aryl is as defined above or
- (28) O(CH₂)_naryl, aryl as defined above;

R³ can also be any of the following when R⁴ is absent:

- (29) oxo,
- 20 (30) =CH-(C₁-C₆)-alkyl, wherein alkyl is as defined above,
- (31) =CH-(C₂-C₆)-alkenyl, wherein alkenyl is as defined above,
- (32) =CH-aryl, wherein aryl is as defined above, or
- (33) =CH₂;

25 R⁸ and R⁹ are independently selected from the group consisting of:

- (1) hydrogen,
- (2) [(C=O)O_R]_Saryl, wherein aryl is as defined above,
- (3) [(C=O)O_R]_S(C₂-C₆)-alkenyl, wherein alkenyl is as defined above,

- (4) $[(C=O)O_r]_s(C_1-C_8)\text{-alkyl}$, wherein alkyl is as defined above,
- (5) $(C=O)_rS(O)_n(C_1-C_8)\text{-alkyl}$, wherein alkyl is as defined above,
- (6) $(C=O)_rS(O)_n\text{-aryl}$, wherein aryl is as defined above, and
- (7) heterocyclyl, wherein heterocyclyl is defined above;

5

R¹⁰ is:

- (1) hydrogen,
- (2) $[(C=O)O_r]_s\text{aryl}$, wherein aryl is as defined above, or
- (3) $[(C=O)O_r]_s(C_1-C_6)\text{-alkyl}$, wherein alkyl is as defined above.

10

2. The Compound of Claim 1, wherein:

R³ and R⁴ are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (2) hydroxy,
- 15 (3) $HO(C_1-C_6)\text{-alkyloxy}$,
- (4) $(C_1-C_4)\text{-perfluoroalkyl}$,
- (5) $O(CO)CCl_3$,
- (6) $(C_1-C_6)\text{-alkyl-S(O)}_n\text{-}$,
- (7) $\text{phenyl-(CH}_2)_r\text{-S(O)}_n\text{-}$,
- 20 (8) cyano,
- (9) nitro,
- (10) CO_2H ,
- (11) $CO(C_1-C_6)\text{-alkyl}$,
- (12) $CO_2(C_1-C_6)\text{-alkyl}$,
- 25 (13) $CONR^8R^9$,
- (14) NR^8R^9 ,
- (15) $O(CO)NR^8R^9$,
- (16) azido,
- (17) $NR^8(CO)NR^8R^9$,

- (18) hydrogen,
- (19) (C₁-C₆)-alkyl, wherein alkyl includes cyclic as well as acyclic groups and is unsubstituted or substituted with one or two of the substituents selected from the group consisting of:
- 5
- (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (b) hydroxy,
- (c) oxo,
- (d) O[(C=O)O]_n(C₁-C₆)-alkyl,
- 10
- (e) (C₁-C₆)-alkyl-S(O)_n-,
- (f) aryl-(C₁-C₆)-alkyloxy,
- (g) cyano,
- (h) nitro,
- (i) vinyl,
- 15
- (j) NR⁸R⁹,
- (k) O(CO)NR⁸R⁹,
- (l) CHO,
- (m) CO₂H,
- (n) CO(C₁-C₆)-alkyl,
- 20
- (o) CO₂(C₁-C₆)-alkyl, wherein alkyl may be substituted with phenyl,
- (p) CONR⁸R⁹,
- (q) aryl, wherein aryl is defined as phenyl or naphthyl, unsubstituted or substituted with one, two or three of the substituents selected from the group consisting of:
- 25
- (a') halo, as defined above,
- (b') hydroxy,
- (c') (C₁-C₆)-alkyl,
- (d') (C₁-C₄)-perfluoroalkyl,

- 5 (e') (C₂-C₆)-alkenyl,
(f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
(i') phenyl,
(j') phenoxy,
(k') cyano,
(l') nitro,
(m') CO₂H,
10 (n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
(q') NR⁸R⁹, and
15 (r) heteroaryl, wherein heteroaryl is defined as an
unsubstituted, monosubstituted, or disubstituted five
or six membered aromatic heterocycle containing
from 1 to 3 heteroatoms selected from the group
consisting of O, N and S and wherein the substituents
are members selected from the group consisting of:
20 (a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
(d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
25 (f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n-,
(i') phenyl,
(j') phenoxy,

- 5 (k') cyano,
(l') nitro,
(m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,
(p') CONR⁸R⁹,
(q') NR⁸R⁹, and
(r') fused benzo or pyridyl group,
- 10 (s) heterocyclyl, wherein heterocyclyl is defined as a 3 to 7 atom cyclic, non-aromatic substituent containing from 1 to 3 heteroatoms selected from the group consisting of O, N, and S, said heterocycle being unsubstituted or substituted with one, two or three substituents selected from the group consisting of:
- 15 (a') halo, as defined above,
(b') hydroxy,
(c') (C₁-C₆)-alkyl,
(d') (C₁-C₄)-perfluoroalkyl,
(e') (C₂-C₆)-alkenyl,
20 (f') (C₂-C₆)-alkynyl,
(g') (C₁-C₆)-alkyloxy,
(h') (C₁-C₆)-alkyl-S(O)_n,
(i') phenyl,
(j') phenoxy,
25 (k') cyano,
(l') nitro,
(m') CO₂H,
(n') CO(C₁-C₆)-alkyl,
(o') CO₂(C₁-C₆)-alkyl,

- 5
- (p') CONR^8R^9 ,
 (q') NR^8R^9 ,
 (r') $\text{NR}^8\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 (s') oxo,
 (t') fused benzo, and
 (u') fused pyridyl group;
- 10
- (t) benzyl- $\text{S}(\text{O})_n$ -,
 (u) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s(\text{C}_2\text{-C}_6)\text{-alkenyl}$,
 (v) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s\text{aryl}$,
 (w) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s\text{heteroaryl}$,
 (x) $\text{O}(\text{CH}_2)_n\text{heteroaryl}$, or
 (y) $\text{O}(\text{CH}_2)_n\text{aryl}$;
- 15
- (20) $(\text{C}_2\text{-C}_6)\text{-alkenyl}$, wherein alkenyl is unsubstituted or substituted with one or two of the substituents selected from the group consisting of:
- 20
- (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (b) hydroxy,
 (c) oxo,
 (e) $(\text{C}_1\text{-C}_6)\text{-alkyl-S}(\text{O})_n$ -,
 (f) phenyl- $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$,
 (g) cyano,
 (h) nitro,
 (i) NR^8R^9 ,
 (j) CHO ,
 25 (k) CO_2H ,
 (l) $\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 (m) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$,
 (n) CONR^8R^9 ,
 (o) aryl, wherein aryl is as defined above,

- (p) heteroaryl, wherein heteroaryl is as defined above,
 (q) heterocyclyl, wherein heterocyclyl is as defined above,
 (r) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
 (s) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
 5 (t) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
 (u) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above,
 (v) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above, and
 (w) $O(CH_2)_n$ aryl, aryl as defined above;
- 10 (21) (C₂-C₆)-alkynyl, wherein alkynyl is unsubstituted or substituted with one or two of the substituents selected from the group consisting of:
- (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (b) hydroxy,
 15 (c) oxo,
 (d) (C₁-C₆)-alkyloxy,
 (e) (C₁-C₆)-S(O)_n-,
 (f) phenyl-(C₁-C₆)-alkyloxy,
 (g) cyano,
 20 (h) nitro,
 (i) vinyl,
 (j) NR^8R^9 ,
 (k) $NR^8CO(C_1-C_6)$ -alkyl,
 (l) CHO,
 25 (m) CO₂H,
 (n) CO(C₁-C₆)-alkyl,
 (o) CO₂C(C₁-C₆)-alkyl,
 (p) CONR⁸R⁹,
 (q) aryl, wherein aryl is as defined above,

- (r) heteroaryl, wherein heteroaryl is as defined above,
- (s) heterocyclyl, wherein heterocyclyl is as defined above,
- (t) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
- (u) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
- 5 (v) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
- (w) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above
- (x) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above, and
- (y) $O(CH_2)_n$ aryl, aryl as defined above,
- (22) $O[(C=O)O_R]_S(C_1-C_6)$ -alkyl, alkyl as defined above,
- 10 (23) $O[(C=O)O_R]_S(C_2-C_6)$ -alkenyl, as defined above,
- (24) $O[(C=O)O_R]_S$ aryl, aryl as defined above,
- (25) $O[(C=O)O_R]_S$ heteroaryl, heteroaryl as defined above
- (26) $O(CH_2)_n$ heteroaryl, heteroaryl as defined above,
- (27) aryl, wherein aryl is as defined above or
- 15 (28) $O(CH_2)_n$ aryl, aryl as defined above;

R^3 can also be any of the following when R^4 is absent:

- (29) oxo,
- (30) $=CH-(C_1-C_6)$ -alkyl, wherein alkyl is as defined above,
- 20 (31) $=CH-(C_2-C_6)$ -alkenyl, wherein alkenyl is as defined above,
- (32) $=CH$ -aryl, wherein aryl is as defined above, or
- (33) $=CH_2$.

3. The compound of Claim 2, wherein:

25 R^1, R^2, R^5, R^6 and R^7 are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (2) hydroxy,
- (3) (C_1-C_6) -alkyl,
- (4) $HO(C_1-C_6)$ -alkyloxy,

- (5) (C₁-C₄)-perfluoroalkyl,
- (6) (C₂-C₆)-alkenyl,
- (7) O[(C=O)O_r]_s(C₁-C₆)-alkyl, wherein the alkyl may be cyclic
or straight-chained,
- 5 (8) phenyl,
- (9) CO(C₁-C₆)-alkyl,
- (10) CO₂(C₁-C₆)-alkyl,
- (11) CONR⁸R⁹,
- (12) NR⁸R⁹,
- 10 (13) (C₂-C₆)-alkenyloxy,
- (14) benzyloxy,
- (15) hydrogen,
- (16) OCF₃,
- (17) R¹ and R² or R⁶ and R⁷ can an be taken together when on
15 adjacent carbons to form a fused benzo, dihydrofuranyl,
furanyl, pyrrolidyl, dihydropyrrolidyl or 1,3-dioxolan group;

R³ and R⁴ are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
- (2) hydroxy,
- 20 (3) HO(C₁-C₆)-alkyloxy,
- (4) (C₁-C₄)-perfluoroalkyl,
- (5) O(CO)CCl₃,
- (6) (C₁-C₆)-alkyl-S(O)_n-,
- (7) phenyl-(CH₂)_r-S(O)_n-,
- 25 (8) cyano,
- (9) nitro,
- (10) CO₂H,
- (11) CO(C₁-C₆)-alkyl,
- (12) CO₂(C₁-C₆)-alkyl,

- (13) CONR^8R^9 ,
(14) NR^8R^9 ,
(15) $\text{O}(\text{CO})\text{NR}^8\text{R}^9$,
(16) azido,
5 (17) $\text{NR}^8(\text{CO})\text{NR}^8\text{R}^9$,
(18) hydrogen,
(19) $(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl includes cyclic as well as
acyclic groups and is unsubstituted or substituted with one
or two of the substituents selected from the group consisting
10 of:
(a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
(b) hydroxy,
(c) oxo,
(d) $\text{O}[(\text{C}=\text{O})\text{O}_r]_s(\text{C}_1\text{-C}_6)\text{-alkyl}$,
15 (e) aryl- $(\text{C}_1\text{-C}_6)\text{-alkyloxy}$,
(f) NR^8R^9 ,
(g) $\text{O}(\text{CO})\text{NR}^8\text{R}^9$,
(h) CHO ,
(i) CO_2H ,
20 (j) $\text{CO}(\text{C}_1\text{-C}_6)\text{-alkyl}$,
(k) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl may be substituted
with phenyl,
(l) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkenyl}$,
(m) CONR^8R^9 ,
25 (n) aryl, wherein aryl is defined as phenyl or naphthyl,
unsubstituted or substituted with a substituent
selected from the group consisting of:
(a') halo, as defined above,
(b') hydroxy,

- 5 (c') (C₁-C₆)-alkyl,
 (d') (C₁-C₆)-alkyloxy,
 (e') (C₁-C₆)-alkyl-S(O)_n-,
 (f) phenyl,
 (g') phenoxy,
 (h') cyano,
 (i') CO₂H,
 (j') CO(C₁-C₆)-alkyl,
 (k') CO₂(C₁-C₆)-alkyl,
 10 (l') CONR⁸R⁹,
 (m') NR⁸R⁹, and
 (o) benzyl-S(O)_n-,
 (p) O[(C=O)O_r]_s(C₂-C₆)-alkenyl,
 (q) O[(C=O)O_r]_saryl,
 15 (r) O[(C=O)O_r]_sheteroaryl,
 (s) O(CH₂)_nheteroaryl, or
 (t) O(CH₂)_naryl;
- 20 (20) (C₂-C₆)-alkenyl, wherein alkenyl is unsubstituted or substituted with a substituent selected from the group consisting of:
- 25 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (b) hydroxy,
 (c) oxo,
 (d) phenyl-(C₁-C₆)-alkyloxy,
 (e) NR⁸R⁹,
 (f) CHO,
 (g) CO₂H,
 (h) CO(C₁-C₆)-alkyl,

- (i) $\text{CO}_2(\text{C}_1\text{-C}_6)\text{-alkyl}$,
- (j) CONR^8R^9 ,
- (k) aryl, wherein aryl is as defined above,
- (l) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}(\text{C}_1\text{-C}_6)\text{-alkyl}$, alkyl as defined above,
- 5 (m) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}\text{aryl}$, aryl as defined above,
- (n) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}\text{heteroaryl}$, heteroaryl as defined above,
- (o) $\text{O}(\text{CH}_2)_\text{n}\text{heteroaryl}$, heteroaryl as defined above, and
- (p) $\text{O}(\text{CH}_2)_\text{n}\text{aryl}$, aryl as defined above;
- (21) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}(\text{C}_1\text{-C}_6)\text{-alkyl}$, alkyl as defined above,
- 10 (22) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}(\text{C}_2\text{-C}_6)\text{-alkenyl}$, as defined above,
- (23) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}\text{aryl}$, aryl as defined above,
- (24) $\text{O}[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}\text{heteroaryl}$, heteroaryl as defined above
- (25) $\text{O}(\text{CH}_2)_\text{n}\text{heteroaryl}$, heteroaryl as defined above,
- (26) aryl, wherein aryl is as defined above or
- 15 (27) $\text{O}(\text{CH}_2)_\text{n}\text{aryl}$, aryl as defined above;

R^3 can also be any of the following when R^4 is absent:

- (28) oxo,
- (29) $=\text{CH}(\text{C}_1\text{-C}_6)\text{-alkyl}$, wherein alkyl is as defined above,
- 20 (30) $=\text{CH}(\text{C}_2\text{-C}_6)\text{-alkenyl}$, wherein alkenyl is as defined above,
- (31) $=\text{CH-aryl}$, wherein aryl is as defined above, or
- (32) $=\text{CH}_2$;

R^8 and R^9 are independently selected from the group consisting of:

- 25 (1) hydrogen,
- (2) $[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}\text{aryl}$, wherein aryl is as defined above,
- (3) $[(\text{C}=\text{O})\text{O}_\text{R}]_\text{S}(\text{C}_2\text{-C}_8)\text{-alkenyl}$, wherein alkenyl is as defined above, and

- (4) $[(C=O)O_r]_s(C_1-C_8)\text{-alkyl}$, wherein alkyl is as defined above,
and

R¹⁰ is:

- 5 (1) hydrogen, or
 (2) $[(C=O)O_r]_s(C_1-C_6)\text{-alkyl}$, wherein alkyl is as defined above.

4. The compound of Claim 3, wherein:

10 R¹, R², R⁵, R⁶ and R⁷ are independently:

- (1) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (2) hydroxy,
 (3) (C₁-C₃)-alkyl,
 (4) (C₂-C₃)-alkenyl,
15 (5) O(C₁-C₄)-alkyl, wherein the alkyl may be cyclic or straight-
 chained,
 (6) O(CO)CH₃,
 (7) CO(C₁-C₃)-alkyl,
 (8) CO₂(C₁-C₃)-alkyl,
20 (9) hydrogen,
 (10) R¹ and R² or R⁶ and R⁷ can an be taken together when on
 adjacent carbons to form a fused benzo, dihydrofuranyl,
 furanyl, pyrrolidyl, dihydropyrrolidyl or 1,3-dioxolan group;

25 R³ and R⁴ are independently:

- (1) hydrogen,
 (2) (C₁-C₆)-alkyl, wherein alkyl includes cyclic as well as
 acyclic groups and is unsubstituted or substituted with one
 or two of the substituents selected from the group consisting
30 of:

- 5 (a) halo, wherein halo is fluoro, chloro, bromo, or iodo,
 (b) hydroxy,
 (c) oxo,
 (d) $O[(C=O)O_r]_s(C_1-C_6)$ -alkyl, wherein r and s are
 independently 0 or 1,
 (e) $CO_2(C_2-C_3)$ -alkenyl,
 (f) $O[(C=O)O_r]_s(C_1-C_6)$ -alkenyl, wherein r and s are
 independently 0 or 1,
 (g) NR^8R^9 ,
 10 (h) $O(CO)NR^8R^9$,
 (i) CHO,
 (j) CO_2H ,
 (k) $CO(C_1-C_6)$ -alkyl,
 (l) $CO_2(C_1-C_6)$ -alkyl, wherein alkyl may be substituted
 15 with phenyl,
 (m) $CONR^8R^9$,
 (n) aryl, wherein aryl is defined as phenyl, unsubstituted
 or substituted with a substituent selected from the
 group consisting of:
 20 (a') halo, as defined above,
 (b') hydroxy,
 (c') (C_1-C_6) -alkyl, and
 (d') (C_1-C_6) -alkyloxy, and
 (o) $O[(C=O)O_r]_s(C_2-C_6)$ -alkenyl, wherein r and s are
 25 independently 0 or 1,
- (3) (C_2-C_6) -alkenyl,
 (4) aryl, wherein aryl is as defined above or
 (5) $O(CH_2)_n$ aryl, wherein aryl is as defined above;

R³ can also be any of the following when R⁴ is absent:

- (6) =CH-(C₁-C₆)-alkyl, wherein alkyl is as defined above,
- (7) =CH-(C₂-C₆)-alkenyl, wherein alkenyl is as defined above,

5

R⁸ and R⁹ are independently selected from the group consisting of:

- (1) hydrogen,
- (2) [(C=O)O_r]_saryl, wherein aryl is as defined above and r and s are independently 0 or 1,
- (3) (C₂-C₈)-alkenyl, wherein alkenyl is as defined above, and
- (4) (C₁-C₆)-alkyl, wherein alkyl is as defined above, and

10

R¹⁰ is:

- (1) hydrogen, or
- (2) (C=O)(C₁-C₃)-alkyl, wherein alkyl is as defined above.

15

5. A compound which is selected from the group consisting of:

1-(2-methoxyphenyl)-1-oxo-2-aza-(S)-4-i-butyl-4-phenylbutane,

20

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-((3-allyloxycarbonyloxy)propyl)-4-phenylbutane,

1-(2-methoxyphenyl)-1-oxo-2-aza-4,4-diethyl-4-phenylbutane,

25

1-(2,3-dihydrobenzofuran-7-yl)-1-oxo-2-aza-4,4-diethyl-4-(phenyl)-butane, and

1-(2-methoxyphenyl)-1-oxo-2-aza-4-(S)-(3-hydroxypropyl)-4-phenylbutane,

30 or a pharmaceutically acceptable salt thereof.

6. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by Kv1.3 inhibition, which comprises administering the compound of Claim 1 in an amount that is effective at inhibiting Kv1.3.

5

7. The method of treating a condition in a mammal the treatment of which is effected or facilitated by Kv1.3 inhibition, as recited in Claim 6, wherein the condition is selected from the group consisting of: resistance by transplantation of organs or tissue, graft-versus-host
10 diseases brought about by medulla ossium transplantation, rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis, juvenile-onset or recent-onset diabetes mellitus, posterior uveitis, allergic encephalomyelitis, glomerulonephritis, infectious diseases caused by
15 pathogenic microorganisms, inflammatory and hyperproliferative skin diseases, psoriasis, atopic dermatitis, contact dermatitis, eczematous dermatitises, seborrhoeic dermatitis, Lichen planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus
20 erythematosus, acne, Alopecia areata, keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, pollen
25 allergies, reversible obstructive airway disease, bronchial asthma, allergic asthma, intrinsic asthma, extrinsic asthma, dust asthma, chronic or inveterate asthma, late asthma and airway hyper-responsiveness, bronchitis, gastric ulcers, vascular damage caused by ischemic diseases and thrombosis, ischemic bowel diseases,
30 inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and leukotriene B₄-mediated

- diseases, Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease, ulcerative colitis, migraine, rhinitis, eczema, interstitial nephritis, Good-pasture's syndrome, hemolytic-uremic syndrome, diabetic nephropathy, multiple myositis, Guillain-
- 5 Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis, mononeuritis, radiculopathy, hyperthyroidism, Basedow's disease, pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, agranulocytosis, pernicious anemia, megaloblastic anemia,
- 10 anerythroplasia, osteoporosis, sarcoidosis, fibroid lung, idiopathic interstitial pneumonia, dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity, cutaneous T cell lymphoma, arteriosclerosis, atherosclerosis, aortitis syndrome, polyarteritis nodosa, myocardosis, scleroderma, Wegener's granuloma,
- 15 Sjogren's syndrome, adiposis, eosinophilic fascitis, lesions of gingiva, periodontium, alveolar bone, substantia ossea dentis, glomerulonephritis, male pattern alopecia or alopecia senilis by preventing epilation or providing hair germination and/or promoting hair generation and hair growth, muscular dystrophy; Pyoderma and
- 20 Sezary's syndrome, Addison's disease, ischemia-reperfusion injury of organs which occurs upon preservation, transplantation or ischemic disease, endotoxin-shock, pseudomembranous colitis, colitis caused by drug or radiation, ischemic acute renal insufficiency, chronic renal insufficiency, toxinosis caused by lung-oxygen or drugs, lung cancer,
- 25 pulmonary emphysema, cataracta, siderosis, retinitis, pigmentosa, senile macular degeneration, vitreal scarring, corneal alkali burn, dermatitis erythema multiforme, linear IgA ballous dermatitis and cement dermatitis, gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution, aging, carcinogenesis, metastasis of
- 30 carcinoma and hypobaropathy, disease caused by histamine or leukotriene-C₄ release, Behcet's disease, autoimmune hepatitis,

primary biliary cirrhosis sclerosing cholangitis, partial liver resection, acute liver necrosis, necrosis caused by toxin, viral hepatitis, shock, or anoxia, B-virus hepatitis, non-A/non-B hepatitis, cirrhosis, alcoholic cirrhosis, hepatic failure, fulminant hepatic failure, late-onset hepatic failure, "acute-on-chronic" liver failure, augmentation of chemotherapeutic effect, cytomegalovirus infection, HCMV infection, AIDS, cancer, senile dementia, trauma, and chronic bacterial infection.

8. The method as recited in Claim 7, wherein the condition is an autoimmune disease.

9. A method of preventing or treating resistance to transplantation or transplantation rejection of organs or tissues in a patient in need thereof, which comprises administering a therapeutically effective amount of a compound of Claim 1.

10. A method of suppressing the immune system in a subject in need thereof, which comprises administering to the subject an immune suppressing amount of a compound of Claim 1.

11. A pharmaceutical formulation comprising a pharmaceutically acceptable carrier and the compound of Claim 1 or a pharmaceutically acceptable crystal form or hydrate thereof.

12. A pharmaceutical composition made by combining the compound of Claim 1 and a pharmaceutically acceptable carrier.

13. The pharmaceutical formulation of Claim 11, comprising, in addition, a second immunosuppressive agent.

14. A pharmaceutical composition made by combining the compound of Claim 1, a pharmaceutically acceptable carrier, and a second immunosuppressive agent.

5 15. The pharmaceutical formulation as recited in Claim 13 wherein the second immunosuppressive agent is selected from a group consisting of azathioprine, brequinar sodium, deoxyspergualin, mizaribine, mycophenolic acid morpholino ester, cyclosporin, FK-506 and rapamycin.

10

16. The method of Claim 10, which comprises coadministering a second immunosuppressive agent.

17. A method of preventing or treating resistance by
15 transplantation of organs or tissue, graft-versus-host diseases brought about by medulla ossium transplantation, rheumatoid arthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis, juvenile-onset or recent-onset diabetes mellitus, posterior uveitis, allergic
20 encephalomyelitis, glomerulonephritis, infectious diseases caused by pathogenic microorganisms, inflammatory and hyperproliferative skin diseases, psoriasis, atopic dermatitis, contact dermatitis, eczematous dermatitises, seborrhoeis dermatitis, Lichen planus, Pemphigus, bullous pemphigoid, Epidermolysis bullosa, urticaria, angioedemas,
25 vasculitides, erythemas, cutaneous eosinophilias, Lupus erythematosus, acne, Alopecia areata, keratoconjunctivitis, vernal conjunctivitis, uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea, dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer, Scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis, pollen
30 allergies, reversible obstructive airway disease, bronchial asthma,

allergic asthma, intrinsic asthma, extrinsic asthma and dust asthma, chronic or inveterate asthma, late asthma and airway hyper-responsiveness, bronchitis, gastric ulcers, vascular damage caused by ischemic diseases and thrombosis, ischemic bowel diseases,

5 inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated with thermal burns and leukotriene B₄-mediated diseases, Coeliac diseases, proctitis, eosinophilic gastroenteritis, mastocytosis, Crohn's disease, ulcerative colitis, migraine, rhinitis, eczema, interstitial nephritis, Good-pasture's syndrome, hemolytic-

10 uremic syndrome, diabetic nephropathy, multiple myositis, Guillain-Barre syndrome, Meniere's disease, polyneuritis, multiple neuritis, mononeuritis, radiculopathy, hyperthyroidism, Basedow's disease, pure red cell aplasia, aplastic anemia, hypoplastic anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia,

15 agranulocytosis, pernicious anemia, megaloblastic anemia, anerythroplasia, osteoporosis, sarcoidosis, fibroid lung, idiopathic interstitial pneumonia, dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity, cutaneous T cell lymphoma, arteriosclerosis, atherosclerosis, aortitis syndrome,

20 polyarteritis nodosa, myocardosis, scleroderma, Wegener's granuloma, Sjogren's syndrome, adiposis, eosinophilic fascitis, lesions of gingiva, periodontium, alveolar bone, substantia ossea dentis, glomerulonephritis, male pattern alopecia or alopecia senilis by preventing epilation or providing hair germination and/or promoting

25 hair generation and hair growth, muscular dystrophy; Pyoderma and Sezary's syndrome, Addison's disease, ischemia-reperfusion injury of organs which occurs upon preservation, transplantation or ischemic disease, endotoxin-shock, pseudomembranous colitis, colitis caused by drug or radiation, ischemic acute renal insufficiency, chronic renal

30 insufficiency, toxinsosis caused by lung-oxygen or drugs, lung cancer, pulmonary emphysema, cataracta, siderosis, retinitis, pigmentosa, senile

macular degeneration, vitreal scarring, corneal alkali burn, dermatitis erythema multiforme, linear IgA ballous dermatitis and cement dermatitis, gingivitis, periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution, aging, carcinogenesis, metastasis of carcinoma and hypobaropathy, disease caused by histamine or leukotriene-C₄ release, Behcet's disease, autoimmune hepatitis, primary biliary cirrhosis sclerosing cholangitis, partial liver resection, acute liver necrosis, necrosis caused by toxin, viral hepatitis, shock, or anoxia, B-virus hepatitis, non-A/non-B hepatitis, cirrhosis, alcoholic cirrhosis, hepatic failure, fulminant hepatic failure, late-onset hepatic failure, "acute-on-chronic" liver failure, augmentation of chemotherapeutic effect cytomegalovirus infection, HCMV infection, AIDS, cancer, senile dementia, trauma, and chronic bacterial infection which comprises administering a compound of Claim 1.

15

18. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by K_v1.3 inhibition, which comprises administering a pharmaceutical formulation comprising a pharmaceutical carrier and a compound of Claim 1, in an amount that is effective at inhibiting K_v1.3.

20

19. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by K_v1.3 inhibition, comprising coadministering a therapeutically effective amount of a compound of Claim 1, with a second immunosuppressive agent.

25

20. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by K_v1.5 inhibition, which comprises administering a compound of Claim 1 in an amount that is effective at inhibiting K_v1.5.

30

21. A method of treating a condition in a mammal, the treatment of which is effected or facilitated by $K_v1.5$ inhibition, which comprises administering a pharmaceutical formulation comprising
5 a pharmaceutical carrier and a compound of Claim 1, in an amount that is effective at inhibiting $K_v1.5$.

22. A method of preventing or treating cardiac arrhythmias in a mammal, which comprises administering a
10 therapeutically effective amount of a compound of Claim 1.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/25042

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :Please See Extra Sheet.

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE, EAST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AVETISYAN et al. Isoquinoline derivatives, Arm. Khim. Zh., 1981, Vol. 34, No. 12, pages 1007-1010, especially page 1009.	1-5 and 11-15
Y	US 5,618,938 A (EMONDS-ALT et al.) 08 April 1997, columns 63-64.	1-6, 8-16, and 18-22
Y	OTT et al. Tetrahydroisoquino [2,1-d] [1,4] benzodiazepines, synthesis and neuro- pharmacological activity. J. Med. Chem., July 1968, Vol. 11, pages 777-787, especially page 778.	1-5, and 11-15

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

14 JANUARY 2000

Date of mailing of the international search report

10 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

D. Lawrence
SHAILENDRA KUMAR

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/25042

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 7 and 17
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

These claims claim method of treating all the possible existing diseases in the human being, by single class of compounds, thus rendering the method claims unduly broad and unsearchable.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/25042

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (7):

A61K 31/27, 31/165, 31/235, 31/265, 31/335; C07C 229/00, 233/65, 271/02; C07D 307/78, 313/04

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

514/450, 469, 478, 512, 534, 617; 549/346, 467; 558/272;
560/42, 157; 564/177

B. FIELDS SEARCHED
Minimum documentation searched
Classification System: U.S.

514/450, 469, 478, 512, 534, 617; 549/346, 467; 558/272;
560/42, 157; 564/177